



apunts

MEDICINA DE L'ESPORT

www.apunts.org



REVIEW

Therapeutic physical exercise and supplements to treat fibromyalgia[☆]



Alfonso Javier Ibáñez-Vera^{a,*}, José Ramón Alvero-Cruz^b,
Jerónimo Carmelo García-Romero^b

^a *Departamento de Ciencias de la Salud de la Universidad de Jaén, Úbeda, Spain*

^b *Departamento de Fisiología Humana, Histología Humana, Anatomía Patológica y Educación Física y Deportiva de la Universidad de Málaga, Málaga, Spain*

Received 27 April 2017; accepted 26 July 2017

Available online 27 October 2017

KEYWORDS

Fibromyalgia;
Exercise;
Supplements;
Aerobic exercise;
Strength

Abstract

Introduction: The prevalence of fibromyalgia in developed countries is over 2.1%. Nowadays, there is no effective treatment for it and different mitigating treatments such as therapeutic exercise are performed instead. In recent years, supplements have also emerged.

Objectives: To review and update the evidence on treatment of fibromyalgia symptoms with therapeutic physical exercise and supplements. To achieve this, treatment application times and the duration of treatment programs will be considered.

Methods: A bibliographic search was carried out on Pubmed database. As a result, 695 studies on therapeutic exercise and 53 on supplements were found. From these, 18 therapeutic exercise studies and 8 supplements studies were selected. They were all chosen and assessed according to their methodological quality, which was measured using the PEDro scale.

Results: Evidence showed that pain and quality of life are improved by the practice of almost any type of physical activity and the intake of Q10 coenzyme or vitamin D supplements. Sleep quality is enhanced by combination of physical exercise and relaxation. Depression, anxiety and mood are improved by the performance of aerobic, flexibility exercises and aquatic biodance.

Conclusions: Therapeutic physical exercise and supplements may be an interesting alternative or complement when treating some fibromyalgia symptoms.

© 2017 FC Barcelona. Published by Elsevier España, S.L.U. All rights reserved.

[☆] Some of the information in this paper was presented as an oral communication at the FEMEDE Congress, held in Granada, November 24, 2016.

* Corresponding author.

E-mail address: ajibanez@ujaen.es (A.J. Ibáñez-Vera).

PALABRAS CLAVE

Fibromialgia;
Entrenamiento;
Suplementos;
Ejercicio aeróbico;
Fuerza

Ejercicio físico terapéutico y suplementos para el tratamiento de la fibromialgia**Resumen**

Introducción: La prevalencia de fibromialgia en los países desarrollados es superior al 2,1%. Actualmente no existe ningún tratamiento eficaz para combatirla y, en su lugar, se efectúan distintos tratamientos para mitigarla, como el ejercicio terapéutico. En los últimos años también han surgido suplementos.

Objetivo: Revisar y actualizar la evidencia sobre el tratamiento de los síntomas de la fibromialgia con el ejercicio físico terapéutico y suplementos. Para lograrlo se revisan los tiempos de aplicación del tratamiento y la duración de estos programas de tratamiento.

Métodos: Se realizó una búsqueda bibliográfica en la base de datos PubMed. Como resultado se encontraron 695 trabajos sobre ejercicio terapéutico y 53 sobre suplementos. A partir de estos se seleccionaron 18 estudios de ejercicio terapéutico y 8 de suplementos. Todos ellos fueron seleccionados y evaluados en función de su calidad metodológica, que se midió utilizando la escala PEDro.

Resultados: La evidencia mostró que el dolor y la calidad de vida mejoraban con la práctica de casi todo tipo de actividad física y con la ingesta de la coenzima Q10 o suplemento de vitamina D. La calidad del sueño mejora mediante la combinación de ejercicio físico y relajación. La depresión, la ansiedad y el estado de ánimo mejoran con la práctica de ejercicio aeróbico, ejercicios de flexibilidad y biodanza acuática.

Conclusiones: El ejercicio físico terapéutico y los suplementos pueden ser una alternativa o complemento interesantes en el tratamiento de algunos síntomas de la fibromialgia.

© 2017 FC Barcelona. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Fibromyalgia is a syndrome characterized by general and widespread pain as well as depression, digestive disorders, fatigue and sleep disturbance.¹ The World Health Organization also includes the presence of other symptoms such as cephalaea, irritable bowel, painful menstruation, irritability, thermal sensitivity, fatigability and articular rigidity.² This disorder is nowadays one of the most serious health issues in our country, as its prevalence rate in developed world countries is 2.1%.³ The American College of Rheumatology states a diagnosis using nociceptor sensitization at a 4 kg/cm² pressure of 11 points out of 18 (previously stated) with a development period of at least 3 months.^{1,4}

Several studies have tried to clarify its etiology, which still remains unknown. Some of these argue that there is an alteration at the level of the neuroendocrine system. Here, a hyporeactivity is produced because of hypothalamic-pituitary-adrenal axis fatigue. This affects metabolism, stress levels and immunologic system.⁵ A prolonged exposure to stress would render certain glandules unable to produce enough hormones in order to maintain the level of alert. As a result, patients would experience symptoms such as fatigue and muscle pain.⁵ On the other hand, a recent study shows that pain experienced by fibromyalgia patients may be produced by the vasodilation of hypothalamus-controlled meta-arterioles, as these may cause patient's body temperature deregulation as well as higher lactic acid accumulation in soft tissues.⁶ Other hypotheses attribute the cause to local alterations, for

instance the higher incidence of myofascial trigger points in patients with fibromyalgia.⁷

No treatment has nowadays proved to be effective when treating fibromyalgia. As a consequence, we focus on the isolated treatment of the different symptoms, mainly using drugs such as analgesics, opioids and antidepressants.⁸ Nevertheless, pharmacological treatment proves not to be very effective to experience long-term pain relief, diverse side effects may appear and dependency and tolerance risk may arise in patients who have been exposed to a long-term opioid analgesic therapy.⁸ This obviously shows it is necessary to find innovative therapeutic solutions so as to minimize side effects and improve symptoms.⁸ Health expenditure levels destined to fibromyalgia raise every year⁹ and because of this, it is highly advisable to know the possible effects of less expensive new treatments which may prove to be more effective.

Therapeutic exercise is presented as an extraordinarily interesting alternative to fibromyalgia pain treatment in its different approaches.⁸ There is evidence on its utility to treat the different symptoms of fibromyalgia⁸ and the number of studies proving this has notably increased in recent years compared to some years ago,^{10,11} mainly because physical exercise constitutes a safe approach and does not present any side effects. Among the reasons why it may help to attenuate fibromyalgia symptoms, we can also mention that aerobic exercise increases blood irrigation to muscles and as a result, these receive more oxygen and the apparition of trigger points is reduced. Furthermore, doing moderate exercise encourages parasympathetic nervous system activation, which reduces the basal activity of the organism and at the same time favors patient's rest and

sleep. Finally, strength training activates both the immune and the endocrine system.¹²

On the other hand, the increase in the number of studies dealing with the use of supplements to fight with fibromyalgia symptoms has been remarkable in recent years. Several supplements, for instance vitamin D^{13,14} or coenzyme Q10¹⁵⁻¹⁷ seem to produce antioxidant effects, which act at mitochondrial level and on the organism energetic metabolism. This results in an improvement in fatigue, pain and quality of life of fibromyalgia patients.¹⁷

The aim of this study is to know the effects and the effectiveness of several therapeutic exercises as well as dietary and pharmacologic supplements to treat pain, quality of life, depression and sleep quality in fibromyalgia patients by revising already published researches on this field. This will enable us to establish some recommendations on therapeutic physical exercise and supplements for patients according to their symptoms.

Methods

A bibliographic search on the kind of exercises performed to fight fibromyalgia symptoms⁸ was carried out. The available literature on Medline database through Pubmed from November 1, 2010 to November 1, 2015 was reviewed. The inclusion criteria were clinical trials included in studies based on participants suffering from fibromyalgia and published from 2010 in English or in Spanish. The exclusion criteria were articles based on unfinished studies or on researches whose intervention area did not deal with physiotherapy or therapeutic exercise.

The terms employed for this search were "fibromyalgia", "treatment" and "therapy" combined with the Boolean operator "and" and the term "exercise". As a result, 695 studies were found. Inclusion criteria were met by 18 of these studies (Fig. 1).

Furthermore, another search was performed in order to know the utility of supplements to treat fibromyalgia symptoms. Again, the source selected was Medline through Pubmed and this time the keywords were "fibromyalgia" and "supplements", joined by the Boolean operator "and". There were 53 results and only 8 of these studies met the established inclusion criteria (Fig. 2).

In order to validate the methodological quality of the selected studies, the Physiotherapy Evidence Database (PEDro) scale was used. It presents 11 answer items ("yes", "no", "no information"). Only 10 of them were evaluable due to the fact that the first one cannot be counted, as it refers to the external validity of the study. One point is given to "yes" items, whereas no punctuation marks were given to the rest of them. Some authors, for instance Moseley et al.¹⁹ state that obtaining 5 punctuation marks or more means a good methodological quality and low risk of bias.¹⁸

Results

In total, 18 studies about physical exercise were analyzed. Those met the established inclusion criteria. According to PEDro scale, there was a good-excellent methodological quality. Punctuation marks ranged from 6 to 10 points out of a maximum of 10. The average value (standard deviation)




Figure 1 Selection of scientific studies containing the terms "fibromyalgia", "treatment", "therapy" and "exercise".




Figure 2 Selection of scientific studies containing the terms "fibromyalgia" and "supplementation".

was 7.83 (1.54) (Table 1). Concerning supplements, 8 studies were analyzed in total. Their methodological quality was good-excellent in 5 of them and poor in 3. Punctuation marks ranged from 4 to 10 and the average

Table 1 Results of application of methodological-quality scale PEDro to research studies on PE.

Research studies	1	2	3	4	5	6	7	8	9	10	Final score
Sañudo, 2015	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7/10
Gavi, 2014	Yes	No	Yes	No	No	Yes	Yes	Yes	Yes	Yes	7/10
Gianotti, 2014	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	6/10
Latorre, 2013	No	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	6/10
López-Rodríguez, 2013	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes	8/10
Kaleth, 2013	Yes	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	8/10
Sañudo 2012	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10/10
Hooten, 2012	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes	8/10
García-Martínez, 2012	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	6/10
Kayo, 2012	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10/10
López-Rodríguez, 2012	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	6/10
Baptista, 2012	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	9/10
Jones, 2012	Yes	No	Yes	No	No	Yes	Yes	Yes	Yes	Yes	7/10
Liu, 2012	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	9/10
Sañudo 2011	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10/10
Arcos-Carmona, 2011	Yes	No	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	8/10
Fontaine, 2010	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	6/10
Sañudo, 2010	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10/10

1, random allocation; 2, concealed allocation; 3, initial comparability; 4, blinding of subjects; 5, blinding of therapists; 6, blinding of assessors; 7, appropriate follow-up; 8, intention-to-treat analysis; 9, between-group statistical comparison; 10, point and variability measures; PE, physical exercise.

Table 2 Results of application of methodological-quality scale PEDro to research studies on supplementation.

Research studies	1	2	3	4	5	6	7	8	9	10	Final score
Wepner, 2014	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10/10
Miyamae, 2013	No	No	Yes	Yes	No	No	Yes	Yes	Yes	Yes	6/10
Cordero, 2013	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	9/10
Alves, 2013	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	10/10
Cordero, 2011	No	No	No	No	No	No	Yes	Yes	Yes	Yes	4/10
Matthana, 2011	No	No	Yes	No	No	No	Yes	Yes	Yes	Yes	5/10
Naziroglu, 2010	Yes	No	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	8/10
Merchant, 2000	No	No	Yes	No	No	No	Yes	Yes	No	Yes	4/10

1, random allocation; 2, concealed allocation; 3, initial comparability; 4, blinding of subjects; 5, blinding of therapists; 6, blinding of assessors; 7, appropriate follow-up; 8, intention-to-treat analysis; 9, between-group statistical comparison; 10, point and variability measures.

value was 7 ± 2.56 (Table 2). Because of the lack of published data on this topic, lower quality studies were also considered.

These are the kind of physical exercises to treat fibromyalgia which were extracted from the chosen studies: four studies were based on aerobic exercise,^{19–22} four studies were based on strength exercises performed with aerobic exercises,^{23–26} two studies were based on the results of strength exercises compared to those of aerobic exercises or walking,^{27,28} one comparative study of strength and flexibility exercises,²⁹ one study exclusively based on strength exercises, one study about a half an hour increase in daily activity together with habit reeducation,³⁰ two studies based on aquatic biodance,^{31,32} one study based on belly dance, one study based on Tai-Chi and finally, one study based on Qigong³³ (Table 2). Most of these studies present a duration of 4.38 ± 1.94 months, a range of 3–9 months and a sample of 62.5 ± 36.36 subjects (Table 3).

Concerning the studies based on supplements, two of them dealt with coenzyme Q10,^{15–17} two studies focused on vitamin D,^{13,14} one study was based on vitamins C and E joined with exercise performance,³⁴ one study was about creatine³⁵ and finally one study on the *Chlorella pyrenoidosa* algae.³⁶ The duration of all of these studies on supplements was variable. The average duration was 22.14 ± 16.07 weeks. This result shows practically no variation when data are extracted from the two low methodological quality (22.2 ± 17.01 weeks). The average number of participants shows no variation either when the three poor methodological quality studies are excluded (43.4 ± 24.69 participants) compared to 42.5 ± 33.29 participants when all studies are considered (Table 4).

Discussion

According to obtained data in our revision session, the kind of supplements and the exercises which could be beneficial

Table 3 Characteristics of clinical studies consulted to elaborate the revision on physical exercise.

Authors, year of publication	Exercise performed	n	Kind of research study	Duration	Results
Sañudo, 2015	AER 60–65% MHR	28	RCT	45–60 min, 2 w, 6 months	Improves DEP and anxiety
Kaleth, 2013	AER 40–65% MHR vs AER + motivation	170	RCT	Progressive 10–30 min increase, 2–3 w, 9 months	Improves pain in group AER + motivation with no differences in group QOF
García-Martínez, 2012	AER 60 at 85% MHR	26	RCT	60 min, 3 w, 3 months	Improves QOF and MO
Arcos-Carmona, 2011	AER in water and RELAX	56	RCT	30 min water and 30 min of relaxation, 2 w, 2.5 months	Improves QOF, SQ and anxiety
Hooten, 2012	EDU + AER 70% MHR vs EDU + STR	69	RCT	30 min/day in 3 w	Both improve pain in a similar way
Gianotti, 2014	AER + STR	32	RCT	60 min, 2 w, 2.5 months	Improves SQ and QOF
Sañudo, 2012	AER 65–70% MHR + STR	41	RCT	45–60 min, 2 w, 6 months; §=2.5 years	Improves QOF and DEP when compared to initial data. Improvement levels are maintained during follow-up
Sañudo 2011	AER 65–70% MHR + STR	42	RCT	45–60 min, 2 w, 6 months	Improves QOF
Sañudo, 2010	AER 65–70% MHR VS AER 56–70% MHR + STR	64	RCT	45–60 min, 2 w, 6 months	Improves MO and QOF in both tre groups with no differences
Latorre, 2013	STR in water and out of water	72	No RCT	60 min, 3 w, 6 months	Improves MO and QOF
Gavi, 2014	STR vs FLEX	66	RCT	45 min, 2 w, 4 months	STR improves pain, FLEX improves anxiety, both improve QOF with no differences
Kayo, 2012	Walking vs STR	90	RCT	3 w, 4 months, §=3months	Both improve QOF and pain
Fontaine, 2010	30 min increase in daily PA vs EDU	92	RCT	6 sessions of 60 min in 3 months	Improves pain and QOF in non-active subjects
López-Rodríguez, 2013	Biodance vs STE	59	RCT	60 min, 2 w, 3 months	Biodance improves SQ, anxiety, QOF and pain when compared to STE
López-Rodríguez, 2012	Biodance vs STE	31	RCT	60 min, 2 w, 3 months	Biodance improves pain, QOF and DEP
Baptista, 2012	Belly dance	75	RCT	60 min, 2 w, 4 months	Improves pain, QOF and MO
Jones, 2012	Tai-chi vs EDU	98	RCT	90 min, 2 w, 3 months	Tai-chi improves QOF, pain and SQ when compared to a EDU
Liu, 2012	Qigong	14	RCT	15 min, 2/day, 6 w	Improves pain, QOF and SQ

STR, strength training; AER, aerobic exercise; FLEX, flexibility exercises; RELAX, relaxation techniques; MHR, maximum heart rate; PA, physical activity; EDU, patient's education; STE, stretching exercises; QOF, quality of life; MO, mood; DEP, depression; SQ, sleep quality; tre, treatment; w, week; §, follow-up after intervention; RCT, randomized clinical trial; No RCT, non-randomized clinical trial.

Table 4 Characteristics of clinical studies consulted to elaborate the revision on supplementation.

Authors, year of publication	Supplement(s)	n	Kind of research study	Duration	Results
Wepner, 2014	Vit D, 20 ng/mL	30	RCT	49 w	Improves pain; QOF, ANS, DEP do not improve pain
Miyamae, 2013	Co Q10 (Ubiquinol-10)	77	No RCT	28 w	Young FBM improves fatigue
Cordero, 2013	Co Q10, 300 mg/day	20	RCT	40 days	Improves pain, QOF and fatigue
Alves, 2013	Creatine, 20 g/day	28	RCT	16 w	It does not improve pain; QOF, DEP only improve strength
Cordero, 2011	Co Q10, 300 mg/day	5	Cases	9 months	Improves pain and QOF
Matthana, 2011	Vit D, 50 IU/w	100	Cohorts	Until Vit D levels reached 50 ng/mL	Improves pain and QOF
Naziroglu, 2010	Vit C and E + Exercise	62	RCT	12 w	No changes in pain
Merchant, 2000	<i>Chlorella pyrenoidosa</i> , 10 g + 10 mL	18	Cohorts	2 months	Improves pain

Co Q10, coenzyme Q10; Vit, vitamin; RCT, randomized clinical trial; No RCT, non-randomized clinical trial; w, weeks; QOF, quality of life; ANS, anxiety; DEP, depression; FBM, fibromyalgia.

for fibromyalgia patients in order to treat their symptoms are diverse. Because of symptom variability in fibromyalgia, the kind of therapeutic exercise and supplements studied have been grouped according to the symptom which proves to be beneficially influenced.

According to Kaleth et al.,²¹ aerobic training at 40% of maximum heart rate (MHR) in a 10-min period of time and progressively incrementing in nine months until reaching 65% AHR in 30 min seems to improve pain, as this does not progress and the initial AHR remains at 40%. In the same line, Fontaine et al.³⁰ suggest that pain improves in fibromyalgia patients who spend at least 30 min on mild daily physical activity in a 3-months period of time. These studies seem to show that all kinds of increase in physical activity – even if this increase is minimum – result in remarkable effects on pain relief. This finding may be of special importance in the case of individuals who present high pain or disability levels.

Concerning strength training, Gavi et al.²⁹ and Kayo et al.²⁸ have proven remarkable pain relief. Nevertheless, Hooten et al.,²⁷ have stated no difference in pain relief when following strength training or aerobic exercises programs at 70% AHR. As a result, beneficial effects in pain relief of both training programs have been proved with no difference among them. This finding may prove that aerobic training at 70% AHR as well as strength training may be useful at the same level. This allows us to choose the most appropriate kind of training program for each patient.

Other therapeutic exercise activities, for instance aquatic biodance show a significant quality-of-life improvement when compared to the effects of stretching treatments.^{31,32} Concerning Tai-chi,³⁷ Qigong³³ and belly dance,³⁸ they also seem to have a significant quality-of-life improvement when compared to the effects of habitual patient care. Fontaine et al.,³⁹ have reported a quality-of-life improvement when mildly augmenting the duration of

daily physical activity – from 30 to 60 min of daily activities performance.³⁹

With regard to supplements, Alves et al.,³⁵ report that a quantity of 20 g of creatine per day does not improve quality of life. Wepner et al.¹³ do not report any improvement neither when administering 20 mg/mL of vitamin D. However, Matthana¹⁴ argues that the quality of life of patients does improve when administering 50 ng/mL of vitamin D. This difference in findings may be due to bias in the study by Matthana, for instance the lack of control and placebo groups or the administration of an insufficient dose by Wepner. As a consequence, vitamin D cannot be recommended in order to improve patients' quality of life until further reliable data are published. Several studies by Cordero et al.^{16,17} state that the quality of life of 25 fibromyalgia patients did improve when a dose of 300 mg of coenzyme Q10 per day was administered. Similar results were proven by Matthana; nevertheless, in this study vitamin D substituted coenzyme Q10. The quantity administered was 50 ng/mL per patient.¹⁴

Considering the information extracted from all the studies previously mentioned, it can be stated that any increase in physical activity positively influences quality of life. Strength training and aerobic exercise programs at 45–85% MHR as well as the combination of these or the performance of activities such as aquatic biodance, Tai-chi, Qigong or belly dance improve quality of life in a major extent than education or habitual patient care. Stretching treatments also improve quality of life, although their effect is less important than that experienced when performing strength training or aquatic biodance. In addition, the intake of coenzyme Q10 and vitamin D may also have beneficial effects; however, further studies on the combination of these with other therapies are required.

Therapeutic physical exercise is also remarkably influential when dealing with depression, anxiety and mood.

Several studies performed by Sañudo et al. suggest that aerobic exercise performed in 45–60 min sessions, 2 days per week in a 6-months period of time significantly improves depression,^{24,40} anxiety⁴⁰ and mood.²⁶ García-Martínez et al.²² reach the same conclusion concerning mood and Arcos-Carmona et al. concerning anxiety²⁰ when similar exercising patterns are followed. Sañudo et al. do not find any differences in the benefits of aerobic exercise when strength training programs are added.²⁶ This fact seems to prove that aerobic exercise is more useful than strength training to improve mood. Gavi et al.²⁹ found similar results. In this study, anxiety decreased in a higher extent in patients who participated in flexibility exercises when compared to results obtained with the performance of strength training exercises.

Meanwhile, studies carried out by López-Rodríguez et al.,^{31,32} show that aquatic biodance improves anxiety³² and depression³¹ in fibromyalgia patients. Belly dance also improves mood when practiced twice a week during several months.³⁸

No studies which improve mood in fibromyalgia patients using supplements have been found. At the same time, no changes when using vitamin D have been found by Wepner et al.¹³ The same occurs when a daily supplements of 20 g of creatine is administered.³⁵

Sleep quality is a strategic aspect in any patient who suffers from chronic pain. The same happens when fibromyalgia chronic pain is concerned.⁴¹ Tang et al. show a relationship between sleep quality and a higher capacity to do physical exercise in fibromyalgia patients.⁴² As a result, sleep quality is considered as an important aspect concerning treatment.

Arcos-Carmona et al.²⁰ report that performing aerobic exercises in water followed by relaxing exercises in sessions of 60 min, twice a week in a period of time of two months and a half significantly improves sleep quality in fibromyalgia patients. A similar result was found by Giannotti et al.,²³ who employed the performance of both aerobic and strength exercises in sessions of an hour, twice a week in a period of time of 10 weeks.

Practicing activities such as aquatic biodance³² and Tai-chi³⁷ twice a week in a two months period of time show significant benefits on fibromyalgia patients' sleep quality.^{32,37} Liu et al.,³³ show very similar results. In this study, a reduced group do Qigong in 15-min sessions, twice a day and during six weeks.³³

When considering data included in the studies consulted, it can be concluded that the influence of supplements on sleep quality still remains unknown due to the fact that no study dealing with this topic has been published.

Taking into consideration all the information mentioned above, the following recommendations for treating fibromyalgia symptoms by performing therapeutic physical exercise can be established:

- Pain: any physical activity which progressively increases the amount of work and which is regularly practiced in 45–60 min sessions, twice or three times a week: strength training, aerobic exercise, aqua exercises, dance activities, Tai-chi, Qigong, walking, etc.
- Quality of life: aerobic exercise performed in 60-min sessions, three times a week and activities which require

dynamic mobilization of the whole body – for instance, dance activities, Tai-chi or Qigong – twice a week.

- Depression, anxiety and mood: regular practice or aerobic and flexibility exercises as well as dance activities in 45–60 min sessions, twice a week.
- Sleep quality: it is advisable to regularly combine aerobic exercises with relaxation or strength training in 60-min sessions, twice a week. Activities such as aquatic biodance, Tai-chi or Qigong can help when practiced twice a week in sessions of 60 min.

It is generally recommended that fibromyalgia patients who present all the symptoms mentioned above do aerobic exercise combined with a progressive-load strength training program in 60-min sessions, three times a week. It is also advisable that they spend two hours per week on dance activities, Tai-chi, Qigong or similar. The intake of 300 mg/day of coenzyme Q10 or maintaining more than 50 ng/mL of vitamin D levels is also recommended.

The lack of research on supplements to treat fibromyalgia should be mentioned as a limitation of this study. In addition, some information may have been missed due to the fact that only the works on therapeutic physical exercise which have been published from 2010 to 2016 have been considered.

In future research, impact of fatigue should be considered to prescribe therapeutic exercise in fibromyalgia patients. Validated tools as the Fatigue Impact Scale⁴³ gives measures that allow to identify subjects who could need an individual physical exercise program whose load would be based on their results in a Maximum Ergometric Test.

Conclusion

- Pain experimented by fibromyalgia patients improve when any physical activity is increased or when this is maintained.
- Fibromyalgia patients' quality of life especially improves with regular performance of therapeutic exercises which require a dynamic mobilization of the patient's body, regardless of load.
- Depression, anxiety and mood improve in fibromyalgia patients when they do regular aerobic and flexibility exercises or activities such as aquatic biodance or belly dance.
- Quality of sleep also improves when doing regular aerobic exercises combined with relaxation, strength training programs and aquatic biodance, Tai-chi or Qigong.
- Supplements may be an interesting therapeutic alternative because of its effectiveness in pain relief and quality of life improvement, however much more studies are needed.

Funding

No scholarships or grants have been received for this work.

Conflict of interests

Authors declare that they don't have any conflict of interests.

References

- Wolfe F, Smythe HA, Yunus MB, Bennet RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia Report of the Multicenter Criteria Committee. *Arthritis Rheum.* 1990;33:160–72.
- Eich W, Häuser W, Friedel E, Klement A, Herrmann M, Petzke F, et al. Definition, classification and diagnosis of fibromyalgia syndrome. *Schmerz.* 2008;22:255–66.
- Wolfe F, Brähler E, Hinz A, Häuser W. Fibromyalgia prevalence, somatic symptom reporting, and the dimensionality of polysymptomatic distress: results from a survey of the general population. *Arthritis Care Res.* 2013;65:777–85.
- Escolar Martín JM, Durán Barbosa R. Fisiopatología de la fibromialgia: alteraciones a nivel cerebral y muscular. *Fisioterapia.* 2011;33:173–82.
- Sommer C, Häuser W, Gerhold K, Joraschky P, Petzke F, Tölle T, et al. Etiology and pathophysiology of fibromyalgia syndrome and chronic widespread pain. *Schmerz.* 2008;22:267–82.
- Albrecht PJ, Hou Q, Argoff CE, Storey JR, Wymer JP, Rice FL, et al. Excessive peptidergic sensory innervation of cutaneous arteriole-venule shunts (AVS) in the palmar glabrous skin of fibromyalgia patients: implications for widespread deep tissue pain and fatigue. *Pain Med.* 2013;14:895–915.
- Alonso-Blanco C, Fernández-de-las-Peñas C, Morales-Cabezas M, et al. Multiple active myofascial trigger points reproduce the overall spontaneous pain pattern in women with fibromyalgia and are related to widespread mechanical hypersensitivity. *Clin J Pain.* 2011;27:405–13.
- García DÁ, Martínez-Nicolas I, Saturno-Hernández PJ. Abordaje clínico de la fibromialgia: síntesis de recomendaciones basadas en la evidencia, una revisión sistemática. *Reumatol Clín.* 2016;65–71.
- Marsico A, Atzeni F, Piroddi A, Cazzola M, Stisi S, Sarzi-Puttini P, et al. Costs of pain in rheumatology. *Reumatismo.* 2014;66:103–7.
- Winkelmann A, Häuser W, Friedel E, Moog-Egan M, Seeger D, Settan M, et al. Physiotherapy and physical therapies for fibromyalgia syndrome. *Der Schmerz.* 2012;26:276–86.
- Ablin J, Fitzcharles M-A, Buskila D, Shir Y, Sommer C, Häuser W, et al. Treatment of fibromyalgia syndrome: recommendations of recent evidence-based interdisciplinary guidelines with special emphasis on complementary and alternative therapies. *Evid Based Complement Alternat Med.* 2013;2013:485272.
- Busch AJ, Barber KR, Overend TJ, Peloso PM, Schachter CL. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev.* 2008;4.
- Wepner F, Scheuer R, Schuetz-Wieser B, Machacek P, Pieler-Bruha E. Effects of vitamin D on patients with fibromyalgia syndrome: a randomized placebo-controlled trial. *Pain.* 2014;155:261–8.
- Matthana MH. The relation between vitamin D deficiency and fibromyalgia syndrome in women. *Saudi Med J.* 2011;32:925–9.
- Miyamae T, Seki M, Naga T, Uchino S, Asazuma H, Yoshida T, et al. Increased oxidative stress and coenzyme Q10 deficiency in juvenile fibromyalgia: amelioration of hypercholesterolemia and fatigue by ubiquinol-10 supplementation. *Redox Rep.* 2013;18:12–9.
- Cordero MD, Alcocer-Gómez E, de Miguel M, Culic O, Carrión AM, Álvarez-Suárez JM, et al. Can coenzyme q10 improve clinical and molecular parameters in fibromyalgia? *Antioxid Redox Signal.* 2013;19:1356–61.
- Cordero MD, Alcocer-Gómez E, De Miguel M, Cano-García FJ, Luque CM, Fernández-Riejo P, et al. Coenzyme Q10: a novel therapeutic approach for fibromyalgia? Case series with 5 patients. *Mitochondrion.* 2011;11:623–5.
- Moseley AM, Herbert RD, Sherrington C, Maher CG. Evidence for physiotherapy practice: a survey of the Physiotherapy Evidence Database (PEDro). *Aust J Physiother.* 2002;48:43–9.
- Del Pozo-Cruz J, Alfonso-Rosa RM, Castillo-Cuerva A, Sañudo B, Nolan P, Del Pozo-Cruz B. Depression symptoms are associated with key health outcomes in women with fibromyalgia: a cross-sectional study. *Int J Rheum Dis.* 2015:1–11.
- Arcos-Carmona IM, Castro-Sánchez AM, Matarán-Peñarrocha GA, Gutiérrez-Rubio AB, Ramos-González E, Moreno-Lorenzo C. Effects of aerobic exercise program and relaxation techniques on anxiety, quality of sleep, depression, and quality of life in patients with fibromyalgia: a randomized controlled trial. *Med Clin.* 2011;137:398–401.
- Kaleth AS, Saha CK, Jensen MP, Slaven JE, Ang DC. Effect of moderate to vigorous physical activity on long-term clinical outcomes and pain severity in fibromyalgia. *Arthritis Care Res.* 2013;65:1211–8.
- García-Martínez AM, De Paz JA, Márquez S. Effects of an exercise programme on self-esteem, self-concept and quality of life in women with fibromyalgia: a randomized controlled trial. *Rheumatol Int.* 2012;32:1869–76.
- Giannotti E, Koutsikos K, Pigatto M, Rampudda ME, Doria A, Masiero S. Medium-/long-term effects of a specific exercise protocol combined with patient education on spine mobility, chronic fatigue, pain aerobic fitness and level of disability in fibromyalgia. *Biomed Res Int.* 2014;2014:1–9.
- Sañudo B, Carrasco L, de Hoyo M, McVeigh JG. Effects of exercise training and detraining in patients with fibromyalgia syndrome: a 3-yr longitudinal study. *Am J Phys Med Rehabil.* 2012;91:561–9.
- Sañudo B, Galiano D, Carrasco L, de Hoyo M, McVeigh JG. Effects of a prolonged exercise program on key health outcomes in women with fibromyalgia: a randomized controlled trial. *J Rehabil Med.* 2011;43:521–6.
- Sañudo B, Galiano D, Carrasco L, Blagojevic M, de Hoyo M, Saxton J. Aerobic exercise versus combined exercise therapy in women with fibromyalgia syndrome: a randomized controlled trial. *Arch Phys Med Rehabil.* 2010;91:1838–43.
- Hooten WM, Qu W, Townsend CO, Judd JW. Effects of strength vs aerobic exercise on pain severity in adults with fibromyalgia: a randomized equivalence trial. *Pain.* 2012;153:915–23.
- Kayo AH, Peccin MS, Sanches CM, Trevisani VF. Effectiveness of physical activity in reducing pain in patients with fibromyalgia: a blinded randomized clinical trial. *Rheumatol Int.* 2012;32:2285–92.
- Gavi MBRO, Vassalo DV, Amaral FT, Macedo DC, Gava PL, Dantas EM, et al. Strengthening exercises improve symptoms and quality of life but do not change autonomic modulation in fibromyalgia: a randomized clinical trial. *PLOS ONE.* 2014;9:e90767.
- Fontaine KR, Conn L, Clauw DJ. Effects of lifestyle physical activity on perceived symptoms and physical function in adults with fibromyalgia: results of a randomized trial. *Arthritis Res Ther.* 2010;12:R55.
- López-Rodríguez MM, Castro-Sánchez AM, Fernández-Martínez M, Matarán-Peñarrocha GA, Rodríguez-Ferrer ME. Comparación entre biodanza en medio acuático y stretching en la mejora de la calidad de vida y dolor en los pacientes con fibromialgia. *Atención Primaria.* 2012;44:641–9.
- López-Rodríguez MM, Fernández-Martínez M, Matarán-Peñarrocha GA, Rodríguez-Ferrer ME, Granados-Gámez G, Aguilar-Ferrándiz E. Efectividad de la biodanza acuática sobre

- la calidad del sueño, la ansiedad y otros síntomas en pacientes con fibromialgia. *Med Clin*. 2013;141:471–8.
33. Liu W, Zahner L, Cornell M, Le T, Le T, Ratner J, Wang Y, et al. Benefit of Qigong exercise in patients with fibromyalgia: a pilot study. *Int J Neurosci*. 2012;122:657–64.
 34. Nazıroğlu M, Akkuş S, Soyupek F, Yalman K, Çelic Ö, Eris S, et al. Vitamins C and E treatment combined with exercise modulates oxidative stress markers in blood of patients with fibromyalgia: a controlled clinical pilot study. *Stress*. 2010;13:498–505.
 35. Alves C, Santiago B, Lima F, Otaduy MC, Calich AL, Tritto AC, et al. Creatine supplementation in fibromyalgia: a double-blind, randomized, placebo-controlled trial. *Arthritis Care Res*. 2013;65:1449–59.
 36. Merchant RE, Carmack CA, Wise CM. Nutritional supplementation with *Chlorella pyrenoidosa* for patients with fibromyalgia syndrome: a pilot study. *Phyther Res*. 2000;14:167–73.
 37. Jones KD, Sherman CA, Mist SD, Carson JW, Bennet RM, Li F. A randomized controlled trial of 8-form Tai chi improves symptoms and functional mobility in fibromyalgia patients. *Clin Rheumatol*. 2012;31:1205–14.
 38. Baptista AS, Villela AL, Jones A, Natour J. Effectiveness of dance in patients with fibromyalgia: a randomized, single-blind, controlled study. *Clin Exp Rheumatol*. 2012;30 Suppl. 74:18–23.
 39. Fontaine KR, Conn L, Clauw DJ. Effects of lifestyle physical activity in adults with fibromyalgia: results at follow-up. *J Clin Rheumatol*. 2011;17:64–8.
 40. Sañudo B, Carrasco L, de Hoyo M, Figueroa A, Saxton JM. Vagal modulation and symptomatology following a 6-month aerobic exercise program for women with fibromyalgia. *Clin Exp Rheumatol*. 2015;33 Suppl. 88:S41–5.
 41. Diaz-Piedra C, Catena A, Sánchez AI, Miró E, Martínez MP, Buelacasa G. Sleep disturbances in fibromyalgia syndrome: the role of clinical and polysomnographic variables explaining poor sleep quality in patients. *Sleep Med*. 2015;16:917–25.
 42. Tang NKY, Sanborn AN. Better quality sleep promotes daytime physical activity in patients with chronic pain? A multilevel analysis of the within-person relationship. *PLOS ONE*. 2014;9.
 43. Shahid A, Wilkinson S, Marcu S. Fatigue Impact scale. In: *Stop, that and one hundred other sleep scales*. 1st ed. New York: Springer; 2011.