



EDITORIAL

Isokinetic values in closed kinetic chain in young adults



KEYWORDS

Dynamometer;
Muscle strength;
Exercise movement
techniques;
Robotics;
Muscle power output;
Functional movement

Abstract

Objective: To describe the values of isokinetic strength in closed kinetic chain (CKC) in young adults and their reproducibility.

Methods: Test and re-test in isokinetic leg press, unilateral mode at 0.1 m/s and 0.3 m/s, 2 sets of 10 repetitions at each speed was performed.

Results: 30 men and 22 women were included. The isokinetic values show differences between men and women. The intraclass correlation coefficient was >0.88 ($p < 0.001$) except in flexors (women), 0.74.

Conclusion: The CKC isokinetic test is valid, reproducible, and safe.

© 2024 CONSELL CATALÀ DE L'ESPORT. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Isokinetics can be defined as a robotic system to evaluate qualitative and quantitative data on muscular effort.¹ It is composed of a constant speed and a variable resistance throughout the range of motion.² The closed kinetic chain (CKC) has been described as “functional” as it is useful in tasks such as standing, sitting, using a ladder, etc.³ Dvir² mentions that the CKC requires multi-joint effort, involves less anterior translation of the tibia and less stress on the anterior cruciate ligament. Davies⁴ adds that the CKC can mask certain localized deficiencies because it compensates with the effort of the other joints.

The CKC is important for movements such as squatting, lunging, jumping, and running, so it must be studied, and analyzed and establish balances that translate for specific tasks.³ We present a cross-sectional study with the aim of describing the values of isokinetic strength in CKC in young adults, the balance between both sides, the flexor/extensor ratio, and their reproducibility.

Methods

Young adults of both genders, without lower limbs or chronic known pathology, were included; a clinical history was taken

to determine the absence of existing or recent joint disease and cardiovascular risk, and informed consent was signed.

Leg Press test was carried out with a Con-Trex LP isokinetic dynamometer, with the following parameters:

Seat inclination 80°, knee extension –20°, hip and knee flexion 90°. Test and re-test in unilateral mode at 0.1 m/s and 0.3 m/s, 2 sets of 10 repetitions at each speed. Warming up and cooling down was performed with continuous passive movement. The values of force were obtained in Newtons (N), muscle power output in Watt (W), differences between both sides, and flexor/extensor ratio.

Data analysis was performed using SPSS software version 24.0; The difference between both sides was analyzed with paired t and reproducibility with intraclass correlation coefficient.

Results

52 people were included, 30 men (average age 30.8 ± 5.3 years, 76.8 ± 10.5 kg weight, 1.72 ± 0.65 m height) and 22 women (average age 29 ± 4.1 years, 59.4 ± 9.6 kg weight, 1.61 ± 0.55 height), who were analyzed separately.

The isokinetic values show a difference of almost 600 N between men and women at 0.1 m/s and 500 N at 0.3 for the extensor chain, with differences between right and left sides of 1% in all cases; and a difference of almost 100 N in the

Abbreviations: CKC, closed kinetic chain

<https://doi.org/10.1016/j.apunsm.2024.100444>

2666-5069/© 2024 CONSELL CATALÀ DE L'ESPORT. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1 Isokinetic values in both groups.

	Men						Women					
	0.1		p	0.3		p	0.1		p	0.3		p
	Average	SD		Average	SD		Average	SD		Average	SD	
N ext dom	1947.2	361.9	0.3	1680.7	290.1	0.13	1346.7	298.6	0.19	1122.8	280.3	0.25
N ext no dom	1963.4	342.8		1650.2	287.3		1367.4	241.4		1108	255.5	
Ratio	0.01			0.01			0.01			0.01		
N flex dom	387.9	66.9	0.09	334.3	53.3	0.25	270.4	43.7	0.12	234.8	25.7	0.02
N flex no dom	398.2	60.7		338.4	53.5		262.5	26.2		222.8	27.8	
Ratio	0.02			0.01			0.02			0.05		
Flex/ext dom	0.2	0.03		0.2	0.03		0.2	0.03		0.21	0.04	
Flex/ext no dom	0.2	0.03		0.2	0.03		0.19	0.02		0.2	0.03	
W ext dom	103.3	33.2		202.2	73		59.9	19.8		112.4	30.6	
W ext no dom	109.1	31		201	78.6		64.4	17.2		119	30.6	
W flex dom	26.6	19		42.3	12.1		15.3	2.7		27.2	5.9	
W flex no dom	27.7	20.6		44.1	13.8		15.4	2.4		27.3	5.4	

SD, standard deviation; N, Newton; Ext, extensors; Flex, flexors; dom: dominant side; No dom, non dominant side; W, Watt.

Table 2 Intraclass correlation in isokinetic tests.

	Men				Women			
	ICC	p	Lim Inf	Lim sup	ICC	p	Lim Inf	Lim sup
N ext 0.1	0.938	<0.001	0.871	0.971	0.962	<0.001	0.905	0.985
N flex 0.1	0.883	<0.001	0.754	0.944	0.809	<0.001	0.518	0.925
N ext 0.3	0.935	<0.001	0.862	0.969	0.968	<0.001	0.918	0.987
N flex 0.3	0.893	<0.001	0.776	0.949	0.741	0.003	0.345	0.897

N, Newton; Ext, extensors; Flex, flexors.

flexors. The flexor/extensor ratio was 20% at both speeds and sexes. These values are shown in [Table 1](#).

Reproducibility was determined with the results of the test and re-test. The intraclass correlation coefficient (ICC) was high for the extensor chains at both speeds and good for the flexors in women ([Table 2](#)).

Discussion

To our knowledge, this is the first research to describe values in young adults with a linear isokinetic dynamometer. Loeza³ used an angular dynamometer, with positioning adapted to CKC, so the values obtained are in degrees per second and not meters per second. Liebensteiner⁵ described values in men and women, however, these were at 0.2 m/s and in eccentric movement, so we have no point of comparison. They found that there were no significant differences in the evaluation between men and women in strength, electromyographic muscle activity, and leg kinematics; in our study, we found a significant difference in both sexes, at both speeds.

Dvir and Müller⁶ describe that in the isokinetic test in CKC, the fixation points are in the hip and foot, but with instability in the knee, however, the forces of the three joints are added to result in a linear vector. In our study, the values obtained have no comparison with those described in a similar population, as it is a physical magnitude different from the angular dynamometry. We can also highlight that the ratio flexor/extensor differs from that obtained with an open chain. Loeza⁷ describes a series of values obtained in

different papers, which are 0.55, unlike what was mentioned by Dvir (0.6)²; in our results, we found a ratio of 0.2 for both groups, at both speeds. This is similar to that described by Loeza³ in young athletes, in the previously mentioned situation of CKC adaptation with an angular dynamometer. This can be understood because the flexor chain (in CKC) does not require the level of force that the extensor chain (antigravity) does.

Reproducibility was high in both groups and at both speeds, since it is not operator-dependent, but rather an exact measurement; similar to that described by other authors regarding the validity of isokinetics. No adverse events occurred during the evaluations.

Our results suggest that the CKC isokinetic test is valid, reproducible, different from another physical evaluation with an angular dynamometer, and safe; and the motor gesture resembles activities of daily life such as walking or climbing stairs; making it an alternative to other strength assessments.

Conflicts of interest

There are no conflicts of interest to declare.

Ethical statements

This protocol was approved by the research, ethics, and bio-safety committees of the Institute of Security and Social

Services of State Workers (ISSSTE) with registration 116.2013, and complies with the Declaration of Helsinki.

Funding

This research received no monetary support or payments during its conduct.

References

- Huesa F, García J, Vargas J. Dinamometría isocinética. *Rehabilitación*. 2005;39(6):288–96.
- Dvir Z. *Isokinetics. Muscle Testing, Interpretation, and Clinical Applications*. 2nd ed, Churchill – Livingstone, 2012.
- Loeza P, Fritzler W, Barrios J. Closed kinetic chain isokinetic values in football players: pilot test. *Arch Med Deporte*. 2017;34(2):66–71.
- Davies GJ, Ellenbecker TS. Chapter 25- application of isokinetics in testing and rehabilitation. *Physical Rehabilitation of the Injured Athlete*. 4th ed Elsevier; 2012, <https://doi.org/10.1016/B978-1-4377-2411-0.00025-3>.
- Liebensteiner MC, Platzer HP, Burtscher M, et al. The effect of gender on the force, muscle activity, and frontal plane knee alignment during maximum eccentric leg-press exercise. *Knee Surg Sports Traumatol Arthrosc*. 2012;20(3):510–6, <https://doi.org/10.1007/s00167-011-1567-0>.
- Dvir Z, Müller S. Multiple-joint isokinetic dynamometry: a critical review. *J Strength Cond Res*. 2020;34(2):587–601, <https://doi.org/10.1519/JSC.0000000000002982>.
- Magaña PL, Solís IGV, Carapia DDF, Morales LEA, Vázquez PIA, González HRQ. Hamstrings/quadriceps ratio in isokinetic tests: are we looking in wrong direction? *Apunts Sports Med*. 2023;58(218).

Pavel Loeza-Magaña^{a,*}, Viridiana Reyes-Bello^a,
Héctor Ricardo Quezada-González^b,
Pedro Iván Arias-Vázquez^c

^a Centro Médico Nacional 20 de Noviembre, av. Félix Cuevas
540. Medicina física y rehabilitación, ciudad de México,
03100, México

^b Sporthabilia, Remedios Valle 19^a, ciudad de México,
México

^c Universidad Juárez Autónoma de Tabasco. Comalcalco,
Tabasco, México

* Corresponding author.

E-mail address: Doctor.pavel@hotmail.com
(P. Loeza-Magaña).

Received 2 February 2024; Accepted 2 April 2024
Available online xxx