

Original Article

Efficacy of a personalized sleep education program and its impact on academic performance in young team sports players¹Ana Merayo^{a,b,c,*}, Oscar Sans^{d,e}, Alex Iranzo^{e,f}, Lluís Capdevila^{g,h}^a Medical Department of Futbol Club Barcelona (FIFA Medical Center of Excellence), Sant Joan Despí, Barcelona, 08970, Spain^b Barça Innovation Hub, Health & Wellness Area, Sant Joan Despí, Barcelona 08970, Spain^c Sports Medicine Unit, Clínic Hospital and Sant Joan de Déu Hospital, Barcelona, 08029, Spain^d Unitat Trastorns del Son. Servei de Neurologia. Hospital Sant Joan de Déu, Esplugues de Llobregat, Barcelona, 08950, Spain^e Faculty of Medicine and Health Sciences, University of Barcelona, Barcelona, 08036, Spain^f Sleep Unit, Neurology Service, Hospital Clínic de Barcelona, IDIBAPS, CIBERNED, 08036, Spain^g Sport Research Institute, Universitat Autònoma de Barcelona, Bellaterra. Barcelona, 08193, Spain^h Laboratory of Sport Psychology, Department of Basic Psychology, Universitat Autònoma de Barcelona, Bellaterra. Barcelona, 08193, Spain

ARTICLE INFO

Keywords:

Academic performance

Athletes

Education program

Sleep quantity

Sleep quality

Sports

Abbreviations: CEIC, scientific research ethics committee

SDSC, sleep disturbance scale in children

SVS, subjective vitality scale

SD, standard deviation

MANOVA, multivariate analysis of variance

ANOVA, Analysis of Variance

CI, Confidence Intervals

SPSS, Statistical Package for the Social Sciences

ABSTRACT

Objective: This study aimed to assess the effectiveness of a sleep education program for young athletes in improving sleep quality, sleep duration, mood, and academic performance.**Design:** Prospective cohort study.**Methods:** A total of 639 athletes (11 % female; mean age: 13.89±3.8 years) from five sports disciplines in a professional club were evaluated before and after a sports season using four specific instruments: (1) sleep diaries to estimate nocturnal sleep duration, (2) the Children's Sleep Disorders Score Scale (SDSC) to assess sleep quality, (3) the Sleep Vitality Scale (SVS) to evaluate mood, and (4) school records to assess academic performance. The sleep education program involved staff, family, and individual sessions, focusing on promoting healthy sleep habits.**Results:** Athletes aged 16–25 years showed an increase in nocturnal sleep duration ($p = 0.002$), whereas those aged 12–15 years experienced a decrease ($p = 0.01$). No changes were observed in the 7–11-year-old group. In terms of sleep quality, significant improvements were observed in the 12–15-year-old ($p < 0.001$) and 16–25-year-old ($p < 0.001$) groups, while the 7–11-year-old group showed a decline ($p = 0.001$). Regarding mood, a significant deterioration was noted in the 7–11-year-old group ($p = 0.008$), with no changes in the older age groups. Academic performance significantly improved across all age groups: 7–11 years ($p = 0.001$), 12–15 years ($p < 0.001$), and 16–25 years ($p = 0.008$).**Conclusions:** The sleep education program effectively improved sleep quality and quantity in the two older age groups (12–25 years). However, it did not yield positive outcomes for the youngest group (7–11 years). All age groups showed a significant increase at the end of the season in academic performance. The sleep education program effectively improved sleep quality, duration, and academic performance in athletes aged 12–25 years. However, it did not yield positive outcomes for athletes aged 7–11 years.

Introduction

There is mounting evidence that sleep plays pivotal role in optimal athletic performance.^{1,2} Adequate sleep optimizes physical, cognitive, and emotional well-being.^{3,4} Studies have demonstrated that partial or total sleep deprivation impairs various aspects of athletic performance,

including physical and neurocognitive functions, mood, body composition, and immune function, among others.^{1,2,5} A growing body of evidence suggests that sleep deprivation may increase the risk of injury in athletes.^{6,7} Adequate sleep quantity and quality has also been identified as a recovery strategy due to its physiological and restorative effects.⁸

With regard to the physical performance of athletes, there is a

¹ All authors have read and approved the final version.

* Corresponding author at. Medical Department of Futbol Club Barcelona (FIFA Medical Center of Excellence), Sant Joan Despí, Barcelona, 08970, Spain.

E-mail addresses: anamerayo@yahoo.es (A. Merayo), osans@hsjdbcn.es (O. Sans), airanzo@clinic.cat (A. Iranzo), lluís.capdevila@uab.cat (L. Capdevila).

growing body of evidence indicating that the sleep deprivation is associated with reduced reaction times in sports such as tennis and basketball,^{9,10} as well as decreased anaerobic power¹⁰ and endurance.^{3,5} Furthermore, insufficient sleep has been linked to negative changes in mood and emotional regulation.¹¹ Factors such as stress, frustration, apathy, irritability, and diminished self-confidence can arise due to lack of sleep.¹² Sleep deprivation has been associated with impaired learning and executive functions, which are crucial for athletes' tactical development during training.^{2,10,13} Other factors, such as demanding training programs, packed schedules, increased travel time, and pre and post-competition anxiety levels, can influence sleep. In the context of sport, it is of paramount importance to maintain an optimal balance between academic pursuits and athletic performance. This is particularly relevant in the case of young athletes, who tend to engage in late-night activities but are required to rise early for academic or training purposes. This can have a detrimental impact on their academic and athletic performance.²

Additionally, there is evidence suggesting that educational interventions can enhance sleep quality and mood in order to improve academic and sport performance. These interventions aim to mitigate or prevent inadequate sleep hygiene by providing tools to manage habits and psychological variables before, during, and after sleep.¹⁴ The interventions focus on reducing hyperactivity, modifying behavior,¹⁵ controlling thoughts,¹⁶ and establishing protocols to enhance sleep.¹⁷ Furthermore, sleep interventions have been demonstrated to be effective in minimizing the negative effects of sleep deprivation and enhancing athletes' overall well-being.¹⁸

The objective of this study was to assess the effectiveness of a personalized sleep education program throughout a sport season in enhancing sleep quality and quantity, psychological well-being, and academic performance among youth players in elite multisport teams.

Methods

Participants

The study included 639 participants (11 % were female), with a mean years of 13.89 ± 3.8 (range, 7 to 25) years. All participants were athletes from various sports disciplines affiliated with a multisport Club including soccer, handball, basketball, futsal, and rink hockey. There was a total of 44 teams among the different categories.

The study was conducted during one academic year and sports season. The study was conducted in accordance with the highest ethical standards and data protection policies, overseen by the club's legal department. All participants were fully informed about the study's details and provided their informed consent to participate. In cases where athletes were under the age of 18 years, legal guardians also signed the informed consent forms. The study protocol (AS-2019-01) was approved by the Scientific Research Ethics Committee (CEIC) of the one University.

Instruments

Sleep diary

The quantity of sleep was evaluated by completing a daily sleep diary each morning upon awakening.¹⁹ This variable will be referred to as "sleep duration". Participants recorded their estimated sleep onset, instances of nocturnal awakenings (if any), and the time and reason for awakening. The hours of sleep were calculated as the sum from bedtime to wake time, subtracting nighttime awakenings and adding the duration of daytime naps. The data were collected over 5 consecutive nights (Monday to Friday, the measure the typical week of the athletes, excluding days without school exams and 4 weekly training sessions), beginning simultaneously with other questionnaires. To determine whether the sleep duration of athletes is optimal or not, we followed the recommendations provided by The National Sleep Foundation (NSF)²⁰

Coaches or captains asked participants to regularly fill out the diary. For statistical analysis, sleep hours were converted to a decimal scale.

Sleep disturbance scale in children (SDSC)

This scale was utilized to assess the sleep quality of participants, with a particular focus on sleep behavior and disturbances.²¹ It encompasses 26 items, grouped into 6 factors, and was completed with the assistance of legal guardians when it was necessary, in accordance with the original study protocol.²¹

Participants and guardians responded to statements pertaining to specific sleep-related behaviors using a Likert-5 scale (1 = Never; 5 = Always; total score range: 26–130). A score of 39 points or above indicates the presence of some degree of sleep disturbance. While the SDSC is typically employed in the assessment of individuals under the age of 18, it was utilized in the evaluation of athletes between the ages of 19 and 25, as there are no discernible clinical scales that differentiate between adolescents and young adults.

Subjective vitality scale (SVS)

This is a validated tool for assessing dynamic well-being and mood.²² The SVS consisted of 7 items that measured enthusiasm, liveliness, and energy levels. Participants responded on a Likert-7 scale ranging from 1 (completely false) to 7 (completely true). The total score is obtained by adding the item scores, with the highest score being 49, indicating greater vitality.

Academic performance assessment

The academic grades for the first (December) and last (June) quarters of the academic year were provided by the participants or their families and recorded for analysis. These grades were obtained from the Spanish educational system. The number of academic subjects and the proportion (%) of subjects passed (grades above 5–10) were documented. Participants were notified of their school grades via letters, in accordance with their signed informed consent.

Procedure

The SDSC and SVS scales, along with the sleep diary were administered at the outset of the season (pre-intervention: PRE) and again at the conclusion of the season (post-intervention: POS), following sleep education sessions in which all players had participated. The instruments were initially completed digitally through an online survey platform by the athletes at the start of the season. The SDSC and SVS took approximately 20 to 30 minutes to complete. All participants completed the instruments during the training sessions. To ensure the collection of more precise and truthful information, legal guardians of players under the age of 12 assisted in completing the instruments alongside the participants. A coach and a member of the research team were present during the instruments implementation process to address any queries. All participants completed the tools on the same day, which was a working day of the week, specifically Monday, and they were away from the weekend match. It was ensured that players filled the tools when they did not have any exams at school. Furthermore, participants were required to maintain a sleep diary for the subsequent 4 days, resulting in a total of 5 entries per participant from Monday to Friday. Each player was required to attend a minimum of 4 sports training sessions during the week. In terms of academic performance, we collected data on the players' grades from the players and their families during the last months of the season. This data provided the percentage of subjects passed by each athlete throughout the academic course.

Sleep education program

The sleep education program was based on a pre-post-intervention design. It was addressed to players and their sports staff, as well as families, as detailed later. Group educational sessions on promoting healthy sleep practices were held for the players' teams, technical staff,

<p>PHASE I: PRE-TEST (MONTHS 1-3)</p> <p>Objective: Baseline assessment and awareness.</p> <ul style="list-style-type: none"> • Participants: Coaches, players, and families • Evaluation: Tools Used <ul style="list-style-type: none"> ○ SDSC and SVS scales ○ Sleep diaries ○ Academic performance evaluations • Actions: <ul style="list-style-type: none"> ○ Provided comprehensive information about the program ○ Explained study objectives and data collection methods ○ Delivered reports on initial sleep duration and quality ○ Offered sleep hygiene improvement recommendations
<p>PHASE II: INTERVENTION PROGRAM (MONTHS 4-8)</p> <p>Objective: Educate and implement sleep hygiene practices</p> <ol style="list-style-type: none"> 1. Sleep Education Sessions for Athletes and Staff <ul style="list-style-type: none"> ○ Participants: Athletes and technical staff (44 teams) ○ Structure: <ul style="list-style-type: none"> ▪ 25-30 min educational session + 10-15 min Q&A ○ Topics Covered: <ul style="list-style-type: none"> ▪ Sleep functions and recommended sleep durations ▪ Sleep quality benefits and routine establishment ▪ Practical tips (sleep schedules, environment, stimulants avoidance, device use, relaxation techniques) ○ Conducted by: Board-certified pediatric sleep specialist 2. Sleep Educational Sessions for Families <ul style="list-style-type: none"> ○ Participants: Athlete families ○ Structure: <ul style="list-style-type: none"> ▪ Similar to athlete session ▪ Focus on phase delay, feeding, and food metabolism ▪ 15-minute Q&A segment ○ Goal: Empower families to support sleep and nutrition needs 3. Individual Counseling Sessions for Athletes <ul style="list-style-type: none"> ○ Frequency: 2 sessions per athlete ○ Focus Areas: <ul style="list-style-type: none"> ▪ Reviewing daily schedules ▪ Teaching relaxation and mindfulness techniques ▪ Implementing personalized sleep strategies ○ Duration: ~1 hour ○ Support Materials: Guides, worksheets, agenda management tools
<p>PHASE III: POST-TEST (MONTHS 9-10)Objective: Evaluation and impact assessment</p> <ul style="list-style-type: none"> • Tools Used: <ul style="list-style-type: none"> ○ SDSC and SVS scales ○ Sleep diaries ○ Academic performance evaluations • Actions: <ul style="list-style-type: none"> ○ Comparative analysis with Phase I data ○ Detailed reports provided to participants and families

Fig. 1.. Sleep education program protocol outline.

and families. In addition, individual counseling sessions were held with each player. A total of 44 educational group sessions were conducted, 1 for each team, from the category with younger players to the category of the affiliate teams for all sports sections. These sessions, attended by players and technical staff (including coaches, doctors, psychologists, and physiotherapists), emphasized essential aspects of sleep hygiene, focusing on promoting healthy sleep practices and strategies to improve sleep quality in young players. All sessions were conducted in person at the Club's. Additionally, group sessions were conducted for the families of all players, regardless of whether they were of legal age or not., with the objective of providing comprehensive knowledge on proper sleep hygiene practices and recommendations. The educational information covered a range of topics, including an overview of sleep and its functions, the physiological changes that occur during sleep, the recommended hours of sleep based on age, and specific recommendations for travel and daily routines.

In addition to the group sessions, 2 sessions were held for each athlete, addressing specific aspects of sleep hygiene relevant to their individual needs. The sessions were tailored to the athletes' schedules, including school and sports schedules and daily routines. All athletes received 2 comprehensive feedback reports, one at the beginning and the other at the end of the season. These report highlighted the quantity and quality of their sleep based on their responses to the sleep diary and the SDSC. The objective of these reports for the athletes was to awareness of their sleep duration and quality, and to raise awareness of their current sleep hygiene.

Characteristics of the educational sleep program

The intervention was executed meticulously in 3 distinct phases, thereby ensuring a comprehensive approach to the promotion of optimal sleep habits in young athletes ((see in Fig.1 the sleep education program protocol outline).

Phase I: pre-test (The initial 3 months of the season). In the initial phase of the study, coaches, players, and all the player's families were provided with comprehensive information and instructions regarding the program's nature and its objectives. Participants were informed about the study's purpose and provided with instructions on how to electronically complete the tools. All participants and their families (in the case of minors) were provided with a detailed report on the results of this initial phase, which included information on their sleep duration and quality, as well as recommendations for improving their current sleep hygiene.

Phase II: intervention program (months 4 to 8 of the intervention). The objective of this phase was to educate athletes on the principles of optimal sleep hygiene, with the aim of improving their sleep quality and duration. The program comprised 3 distinct types of sessions (Fig. 1):

Sleep education session for athletes and staff. These group sessions were conducted for all athletes of the same sport category together with their technical staff (coach, doctor, psychologist and physiotherapist), with a total of 44 teams in different categories. All participants attended a session led by a board-certified pediatrician sleep specialist. This 25–30 mins educational session was followed by a 10–15 mins questions and answers period. A total duration of 35–45 mins was proposed to maintain the maximum attention of the players and to allow for extra time to resolve any doubts that the participants may have on the sleep program. The subject was the same for all groups and the main topics covered included understanding the functions of sleep, recommended sleep times according to age, the benefits of good sleep quality, and recommendations to improve sleep quality and establish effective sleep routines. The professional who conducted the session adapted the vocabulary and examples to the maturational age of each category to facilitate understanding by the study population. The focus was on maintaining to regular sleep schedules, creating a quiet, dark, and cool sleep environment, avoiding stimulants before bedtime, caring for

Table 1.
General characteristics of the sample.

Gender	n	%	
Sport	Men	569	89
	Women	70	11
Sport	n	%	
	Football (men)	277	43.3
	Football (women)	70	11.0
	Basketball	80	12.5
	Futsal	65	10.2
	Hockey	41	6.4
	Handball	106	16.6
Age Categories	n	%	
	7–12 years	153	23.9
	13–15 years	293	45.9
Academic	16–19 years	193	30.2
	n	%	
	Primary School	159	24.8
	Secondary School	286	44.7
	High School	112	17.5
	University	36	5.6
Overall	Other	46	7.1
	639	100 %	

Data are reported as number (n) and percentage (%) of athletes.

mattresses and pillows, limiting electronic device use 2 h before bedtime, avoiding naps exceeding 20 mins, eating a light dinner 2 h before bedtime, and cultivating a calm state before going to sleep.

Sleep educational session for families. A parallel session was conducted exclusively for families, maintaining the same structure of the group session for athletes and staff. The information provided covered technical aspects such as phase delay, feeding, and food metabolism. The session was followed by a 15 mins question-answer segment facilitated by the sleep specialist. The objective was to provide families with information on how to support their athlete's sleep and nutrition needs. It was recommended that athletes and their families implement the recommendations discussed during these sessions over the subsequent weeks.

Individual counseling session for athletes. All athletes engaged in 2 individual counseling sessions, 1 for each age group. The focus was to review their daily schedules, teach relaxation and mindfulness techniques, and implement the recommendations elucidated during the educational session. The topics covered in this study included the reduction of electronic devices usage, avoiding dinner 2 h before bedtime, and maintaining adequate light and temperature in the bedroom. The support materials provided included recommendations, worksheets, sleep and rest guides, and agenda management tools. The sessions were approximately 1h in length, and athletes were permitted to consult with their psychologist at any time during the day. These professionals were also present during the educational sessions addressed to family and staff.

Phase III: post-test (the last two months of the intervention). At the conclusion of the study, participants were evaluated using the SDSC and SVS scales, as well as the sleep diary and academic performance evaluations. This phase served as a follow-up to assess the impact of the intervention. All participants and their families (in the case of minors) received a detailed report on the results of this 3rd phase, including a comparative analysis between results from the first and third phases (Fig. 1)..

Data analysis

The data were presented as means \pm standard deviation (SD) for continuous variables, and participant numbers (n) and percentages (%) for categorical variables. Participants were divided into three groups according to their age, namely group 1 comprising athletes between 7 and 11 years, group 2 between 12 and 15, and group 3 between 16 and 25. A mixed Multivariate Analysis of Variance (MANOVA) with a 3×2

Table 2.
PRE-POS-intervention differences for the variables of the study by age group.

MEASURE	Age Categories	n	PRE Intervention	POS Intervention	Difference (95 % CI)	Cohen's d effect size	p
Sleep Duration (hours)	1 (7–11yo.)	48	9.41 ± 0.12	9.40 ± 0.16	- 0.02 (-0.27 to 0.24)	0.88	0.444
	2 (12–15yo)	150	8.52 ± 0.69	8.31 ± 0.90	- 0.21 (-0.39 to -0.03)	1.11	0.011
	3 (16–26yo.)	72	7.88 ± 0.99	8.20 ± 0.13	0.32 (0.01 to 0.63)	1.32	0.023
	Overall	270	8.51 ± 0.98	8.48 ± 1.18			
Sleep Quality (SDSC score)	1 (7–11yo.)	62	35.82 ± 1.14	43.46 ± 1.28	7.65 (4.50 to 10.79)	12.39	< 0.001
	2 (12–15yo)	180	46.70 ± 0.67	42.7 ± 0.70	-3.96 (-5.97 to -1.94)	13.73	< 0.001
	3 (16–26yo.)	99	51.27 ± 0.95	43.37 ± 1.01	-7.90 (-10.48 to -5.31)	12.97	< 0.001
	Overall	341	46.05 ± 10.32	43.06 ± 10.09			
Mood Assessment (SVS score)	1 (7–11yo.)	6	30.67 ± 2.25	19.67 ± 7.09	-11.00 (-18.94 to -3.06)	7.56	0.008
	2 (12–15yo)	74	25.51 ± 4.92	24.19 ± 5.57	-1.32 (-3.10 to 0.45)	7.67	0.071
	3 (16–26yo.)	54	24.46 ± 4.33	25 ± 4.77	0.54 (-0.92 to 1.99)	5.33	0.231
	Overall		25.32 ± 4.75	24.31 ± 5.40			
Academic Performance (%)	1 (7–11yo.)	153	98.70 ± 4.76	99.80 ± 1.39	1.10 (0.38 to 1.81)	4.46	0.001
	2 (12–15yo)	289	91.63 ± 15.76	96.11 ± 12.27	4.47 (3.09 to 5.85)	11.92	< 0.001
	3 (16–26yo.)	140	77.12 ± 28.76	82.43 ± 27.48	5.31 (1.01 to 9.61)	25.73	0.008
	Overall	582	90.01 ± 19.71	93.79 ± 17.30			

SDSC: Sleep Disturbance Scale in Children. SVC: Subjective Vitality Scale.




Fig. 2. Evolution of sleep hours from the sleep diary between PRE and POS-intervention by age group.
Caption: * ($p < 0.001$: PRE-POS differences with respect to 7–11 years old group. Bars: Mean and SEM).

design (representing 3 age groups * 2 within-subject PRE-POS situations) was used to explore global differences between pre and post-intervention across all study parameters within each age group. A repeated-measures analysis of variance (ANOVA) was conducted within each age level to assess differences between the PRE and POS intervention periods. Mean differences, along with 95 % confidence intervals (CI), and Cohen's d effect sizes were calculated. All statistical analyses were performed using the SPSS statistical package (IBM, SPSS V25.0, Chicago, IL). Results with p-values <0.05 were considered statistically significant.

Results

General characteristics of the sample

Table 1 (General Characteristics of the sample) presents the general characteristics of the sample. It should be noted that only soccer had a contingent of female participants, as detailed in this table. The cohort consisted of 639 athletes (89 % male, 11 % female), with football being the only sport including female participants (11 %). Age distribution revealed three categories: 7–12 years (23.9 %), 13–15 years (45.9 %),




Fig. 3.. Percentage of athletes by age group according to the variation in minutes slept between POS and PRE-intervention, recorded with the Diary (negative values indicate that athletes have slept less in the POS).

Groups: 7–11 years old, 12–15 years old and 16–25 years old. Min=Minutes.

and 16–19 years (30.2 %). Regarding academic background, most athletes were enrolled in secondary school (44.7 %), followed by primary school (24.8 %), high school (17.5 %), and university (5.6 %).

Pre-Post intervention analysis

Table 2 outlines the pre- and post-intervention differences for the study variables across age groups. A mixed MANOVA (3×2) identified significant differences in sleep duration, sleep quality, mood, and academic performance between pre- and post-intervention periods.

The results demonstrated significant variations in sleep hour evolution between PRE and POS interventions across the 3 age groups ($p = 0.006$).

Sleep duration

The analysis of sleep duration, as recorded in the sleep diary, revealed distinct patterns among different age groups. Table 2 shows the duration of sleep in a decimal scale. Specifically, the 7–11-year-old group exhibited no changes between PRE and POST intervention periods. In contrast, the group aged 12–15 experienced a significant reduction in sleep hours (12.6 mins in hexadecimal scale), while the group aged 16–25 showed an increase in sleep duration during the same period (19.2 mines in hexadecimal scale). Fig. 2 illustrates these differences. In particular, the 12–15-year-old group exhibited a significant decrease in sleep hours in comparison to the 7–11-year-old group ($p < 0.001$), which maintained a stable sleep duration. Conversely, the group aged 16–25-year-old group exhibited a significant increase in sleep hours, when compared with the 7–11-year-old group ($p < 0.001$) (see

Table 2 and Fig. 2). Fig. 4 presents the percentage of athletes in each group displaying differences in hours slept between POS and PRE-intervention period, as recorded in the Diary. Differences (POS-PRE) are grouped into 15 mins intervals (a negative value indicates that athletes slept less during POS period).

The analysis of sleep duration, as recorded in the sleep diary, revealed distinct patterns among different age groups. Table 2 shows the duration of sleep in a decimal scale. Specifically, the 7–11-year-old group exhibited no changes between PRE and POST intervention periods. In contrast, the group aged 12–15 experienced a significant reduction in sleep hours (12.6 mins in hexadecimal scale), while the group aged 16–25 showed an increase in sleep duration during the same period (19.2 mines in hexadecimal scale). Fig. 2 illustrates these differences. In particular, the 12–15-year-old group exhibited a significant decrease in sleep hours in comparison to the 7–11-year-old group ($p < 0.001$), which maintained a stable sleep duration. Conversely, the group aged 16–25-year-old group exhibited a significant increase in sleep hours, when compared with the 7–11-year-old group ($p < 0.001$) (see Table 2 and Fig. 2).

The percentage of athletes in each group displaying differences in hours slept between POS and PRE-intervention period is presentment in Fig. 3 presents as recorded in the Diary. Differences (POS-PRE) are grouped into 15-minute intervals (a negative value indicates that athletes slept less during POS period). These findings demonstrate differential effects of the intervention, underscoring the need for tailored strategies to enhance sleep quality and duration, particularly among adolescent athletes.




Fig. 4.. Evolution of SDSC scorings between PRE and POS-intervention by age group. Scores below 39 (red line) indicate higher sleep quality (within the recommended range³⁸).

Caption: * ($p < 0.001$): PRE-POS differences with respect to 7–11 years old group. Bars: Mean and SEM).

Sleep quality

When assessing sleep quality was assessed using the SDSC (higher scores indicating poorer sleep quality), the group aged 7–11 displayed a decline in sleep quality. In contrast, both the 12–15 and 16–25-year-old groups exhibited an improvement in sleep quality. Fig. 4 illustrates this distinct behavior of the 7–11-year-old group, which exhibited an increase in the SDSC score ($p < 0.001$) compared to the improvement shown by the other 2 age-groups ($p < 0.001$). Furthermore, the group aged 16–25 exhibited a more pronounced decline decrease in SDSC score during POS compared to the group aged 12–15 groups (see Table 2 and Fig. 4).

Regarding mood

Regarding mood, as assessed with the SVS (a higher score indicates a better mood), the 7–11-year-old and the 12–15-year-old groups exhibited a decrease in mood whereas the 16–25-year-old group exhibited no difference. Fig. 5 illustrates that the group aged 7–11 exhibited a lower mood score in POS with respect to the PRE ($p = 0.008$), compared to the other two groups (see Table 2 and Fig. 5).

Academic performance

All age groups showed a significant increase at the end of the season in academic performance. Fig. 6 illustrates this change, as measured by the percentage of passed subjects between the PRE and POS periods across different age groups. The entire sample demonstrates an improvement in academic performance during the POS period, increasing from 90 % in the PRE period to 93.79 % in the POS period. However, the 12–15 ($p < 0.001$) and 16–25 ($p = 0.008$) age groups exhibited a more pronounced enhancement in academic performance

during the POS period in comparison to the 7–11 age group ($p = 0.001$) (see Table 2 and Fig. 6). No significant differences were found between genders or among different sports in any of the studied results.

Table 3 (Change in the scores of the main study variables in the POS compared to the PRE-intervention according to the years age group) presents a summary of the changes in study variables from the PRE to the POS-intervention, with POS serving as a reference point. In summary, for children aged 7 to 11, sleep duration remains consistent in both the PRE and POS-intervention periods. However, in this group there is a decrease in sleep quality and mood scores, and an improvement in academic performance. In the 12–15 age group, there was a decrease in sleep duration and mood during the post-intervention period, but an increase in sleep quality and academic performance. Finally, in the 16–25 age group, there was an increase in sleep duration, sleep quality, and academic performance during the post-intervention period, while mood remained consistent.

Discussion

The objective of this study was to evaluate the efficacy of a customized sleep education program carried out during a single season in young athletes. The intervention was designed to improve the quality and quantity of sleep, psychological well-being, and academic performance of 639 athletes who participated in an elite multi-sport team. Overall, the intervention was effective across all age groups, with the 16 to 25-year-old group showing the most significant improvement. The intervention resulted in increased sleep duration, improved sleep quality, and enhanced academic performance. Academic performance improved across the entire sample, and sleep quality also improved in




Fig. 5.. Evolution of mood scorings between PRE and POS-intervention by age group.
Caption: * ($p = 0.008$: PRE>POS for 7–11 years old group. Bars: Mean and SEM).

the intermediate age group of 12 to 15 years. However, the youngest age group (7 to 11 years old) experienced a decline in sleep quality. Mood did not improve in any age group and deteriorated in younger participants.

Sleep duration: The older age group showed the most pronounced increase in sleep duration, affecting approximately 15 % of athletes in this group, resulting in an increase of 19.2 mins in sleep time. This group was the only one averaging <8 h of sleep in the PRE period. In contrast, the intermediate age group experienced a significant decrease in sleep duration, with approximately 13 % of athletes in this group sleeping 12.6 mins less. Most athletes in this group exhibited negative values indicating a reduction in sleep hours during the POS intervention period (see Fig. 3).

This study primarily included soccer players²³, but the results are applicable to other sports, allowing for generalization. Previous research²⁴ focused on soccer players and revealed that they generally slept fewer hours than recommended for non-athlete populations of the same age according to the National Sleep Foundation.²⁵ Notably, only the older age group (16–25 years) exhibited an increase in sleep duration post-intervention, with an average of 8.20 h per night (see Table 2). The average sleep duration was below the recommended levels set forth by the National Sleep Foundation for the 7–11-year and 12–15-year age groups. The latter age group exhibited a reduction in the average duration of sleep, which may be attributed to an increase in mobile phone screen time,²⁶ late-night training schedules,²⁷ and the difficulty of reducing physical and mental arousal after training sessions.

Sleep quality: The SDSC questionnaire was utilized to assess sleep quality, which demonstrated a significant improvement in the 2 older age groups. This resulted in a reduction in SDSC scores. However, the youngest age group (7 to 11 years old) exhibited a decline in sleep

quality post-intervention. Notably, this age group had the highest sleep quality before the intervention, with scores within the recommended range according to the SDSC questionnaire (scores below 39 indicate higher sleep quality²¹). We believe that these results may be related to multifactorial causes: time of the season when it was decided to continue or not in the Club, change of daylight hours with more daily light increase, more competition and tournaments... The intervention appeared to have equalized sleep quality scores among all groups following the intervention. However, all groups scored above the cutoff point of 39 points for poor sleep quality (see Fig. 4). Improving sleep quality is simple and effective, benefiting not only athlete health but also athletic performance.²⁸

In young athletes, the combination of poor sleep quality and insufficient sleep hours indicates a need for specialized attention to improve their sleep-wake habits. Interventions that aim to promote personalized and intensive healthy sleep habits, established at an early age, are crucial for young athletes' lifestyles.²⁵ The effectiveness of the current intervention may have been insufficient for younger participants, highlighting the need for personalized interventions that are tailored to players and families, considering age, particularly for younger athletes,²⁹ equipping them with tools to lower activation, routines that facilitate better daily organization, and thought detection for moments of overthinking. Moreover, older athletes, who benefited more from the intervention, might be more aware of the relationship between recovery behaviors and performance due to their greater mental maturity, underscoring the importance of managing their sleep habits.³⁰

Academic performance: All athletes, regardless of age group, demonstrated an improvement in their pass percentage following the intervention. The improvement was significantly higher in the 2 older age groups, with an approximate 5 % increase in the number of subjects




Fig. 6.. Evolution of percentage of academic passed between PRE and POS-intervention by age group. Caption: * ($p < 0.01$: POS>PRE for all age groups. Bars: Mean and SEM).

Table 3. Change in the scores of the main study variables in the POS compared to the PRE-intervention according to the years age group.

Years Age Group	Quantity Sleep	Quality Sleep	Mood	Academic Results
7-11	=	↓	↓	↑
12-15	↓	↑	↓	↑
16-25	↑	↑	=	↑

↑ Improvement; ↓ Getting worse; = Remains the same.

passed during the POS intervention period compared to PRE intervention period (see Fig. 6). The observed increase in the effectiveness of sleep quantity and quality with the intervention could be related to the progressive academic challenges associated with age. Nevertheless, it is crucial to acknowledge that the 7–11-year-old cohort exhibited a remarkably high pass percentage in PRE (97.8 %), nearly reaching the maximum pass percentage in the POS intervention period (99.8 %) (see Table 2 and Fig. 6). Previous studies have demonstrated that sleep deprivation or poor sleep quality can have a detrimental impact on mental acuity, daily neurocognitive abilities, and academic performance in children and adolescents. Furthermore, it can impair daily

functioning and learning abilities.³¹

Athletes are able to perform well under pressure, even in inadequate conditions. However, it should be noted that constant sleep deprivation can negatively affect performance and that such circumstances can sometimes result in poor performance. It’s evident that continued sleep deprivation cannot sustain consistently good results.²⁴

Educational training: our findings underscore the significance of interventions to enhance sleep quantity and quality among young athletes across the lifespan.²⁴ Adolescents in particular should be encouraged to alter their sleep habits through the incorporating of personalized strategies.³² Sleep hygiene is important for promoting overall health and should be accompanied by proper nutrition, hydration, mood optimization, and stress management. These habits are essential not only in sports but also in personal and academic life. It is of the utmost importance to cultivate self-regulating and responsible behaviors in young athletes. Parents and sports staff should be involved in promoting and reinforcing these habits.^{33,34}

Strengths and limitacions

The study has several strengths. It includes a large sample of young athletes from Futbol Club Barcelona, a Club with a strong tradition of

caring for athletes, promoting invisible training and self-care. Additionally, the study was conducted in a supportive environment. Finally, this study is the first to research and intervene within the same club with such a large number of participants. It provides information on the amount and quality of sleep in a wide and varied sample of athletes, which had not been reported previously.

It should be noted that the study has certain limitations. The study design was based on specific scales and self-report data to collect information about sleep quality and quantity. No objective instruments were used to assess sleep such as the use of mHealth wearable actigraphy and/or polysomnography. The use of actigraphy would have provided data regarding circadian variations between the 3 age groups.³⁵ To control for possible bias, responses from both the group and their parents were included. Another limitation was the duration of the intervention, which was limited to a single season. Extending the intervention period could have revealed additional effects in the study population. Additionally, evaluating sleep data over a limited number of days across 1 season was a limitation. Finally, there is a gender bias as in our study women represented only 11 % of the cohort and all women played soccer.

Practical applications and future research

The results of this study could lead to the development of new practical applications and preventative measures not only in the sports population but also in the general population, including early ages and adults. Educational interventions promoting healthy sleep and lifestyle habits could be implemented as a routine among the athletes and their clubs. In the case of this study, sleep hygiene education training for athletes and the sporting environment at the beginning of each season and the use of individual guidance and follow-up to optimize athletes' rest is implemented. Dissemination and training on sleep pose a significant challenge. It is important to inform and educate athletes. However, the challenge lies in training the educators and ensuring that families, athletes, and coaches/sports staff receive the same teaching regarding the implementation of adequate sleep habits. The researchers of this study aim to investigate a personalized intervention in sleep hygiene to achieve better results, and explore the relationship between the quantity and quality of sleep, school performance, and mood.

Conclusion

In conclusion, the results of our study indicate that:

- The 2 older age groups, particularly the 16 to 25-year-old group, exhibited significant improvements in both sleep quantity and quality following participation in a sleep educational program.
- The entire cohort demonstrated enhanced academic performance while the mood remained stable across all age groups.
- The intervention improved sleep quality for all groups, with the exception of the youngest group, aged 7 to 11 years old, who experienced a decline.
- The results emphasize the importance of continuous efforts to enhance sleep quality and duration in young athletes, with a strong emphasis on tailoring interventions to different age groups to achieve maximum effectiveness.

Formatting of funding sources

This work was supported from Grants PID2019-107473RB-C21, funded by the Spanish Government (MCIN/AEI/ 10.13039/501100011033), and 2021SGR-00806, funded by the Catalanian Government.

Conflicts of interest

The authors declare that they have no competing interest.

Acknowledgments

We extend our gratitude to all FC Barcelona players and staff for their cooperation during the data collection phase. Special thanks to Veronica Tutte and Mario Reyes for their valuable insights and support during the writing of this article.

References

1. Halson SL, Juliff LE. Sleep, sport and the brain. *Prog Brain Res.* 2017;234:165–178. <https://doi.org/10.1016/bs.pbr.2017.06.006>.
2. Malhotra RK. Sleep recovery and performance in sports. *Neurol Clin.* 2017;35(3): 547–557. <https://doi.org/10.1016/j.ncl.2017.03.002>.
3. Simpson NS, Gibbs EL, Matheson GO. Optimizing sleep to maximize performance: implications and recommendations for elite athletes. *Scand J Med Sci Sports.* 2017;27(3):266–274. <https://doi.org/10.1111/sms.12703>.
4. Wheaton AG, Liu Y, Perry GS, Croft JB. Effect of short sleep duration on daily activities-United States, 2005-2008. *Morb Mortal Wkly Rep.* 2011;60:239–242.
5. Fullagar HK, Skorski S, Duffield R, Hammes D, Coutts AJ, Meyer T. Sleep and athletic performance: the effects of sleep loss on exercise performance and physiological and cognitive responses to exercise. *Sports Med.* 2015;45(2):161–186. <https://doi.org/10.1007/s40279-014-0260-0>.
6. Milewski MD, Skaggs DL, Bishop GA, et al. Chronic lack of sleep is associated with increased sports injuries in adolescent athletes. *J Pediatr Orthop.* 2014;34(2): 129–133. <https://doi.org/10.1097/BPO.0000000000000151>.
7. Von Rosen P, Frohm A, Kottorp A, Fridén C, Heijne A. Multiple factors explain injury risk in adolescent elite athletes: applying a biopsychosocial perspective. *Scand J Med Sci Sports.* 2017;27(12):2059–2069. <https://doi.org/10.1111/sms.12855>.
8. Halson SL. Nutrition, sleep and recovery. *Eur J Sport Sci.* 2008;8(2):119–126. <https://doi.org/10.1080/17461390801954794>.
9. Suppiah HT, Low CY, Chia M. Effects of sport-specific training intensity on sleep patterns and psychomotor performance in adolescent athletes. *Pediatr Exerc Sci.* 2016;28(4):588–595. <https://doi.org/10.1123/pes.2015-0205>.
10. Watson AM. Sleep and athletic performance. *Curr Sports Med Rep.* 2017;16(6): 413–418. <https://doi.org/10.1249/JSR.0000000000000418>.
11. Ong AD, Kim S, Young S, Steptoe A. Positive affect and sleep: a systematic review. *Sleep Med Rev.* 2017;35:21–32. <https://doi.org/10.1016/j.smrv.2016.07.006>.
12. Venter RE. Role of sleep in performance and recovery of athletes: a review article. *S Afr J Res Sport Phys Educ Recreat.* 2012;34(1):167–184.
13. Pallesen S, Gundersen HS, Kristoffersen M, Bjorvatn B, Thun E, Harris A. The effects of sleep deprivation on soccer skills. *Percept Mot Skills.* 2017;124(4):812–829. <https://doi.org/10.1177/0031512517707412>.
14. García HBÁ, Correa UJ. Intervención psicológica en trastornos del sueño: una revisión actualizada. *Clínica Contemporánea.* 2020;11(2).
15. Bootzin R. Stimulus control treatment for insomnia. *Proc Annu Conv Am Psychol Assoc.* 1972;7:395–396.
16. Harvey A, Payne S. The management of unwanted pre-sleep thoughts in insomnia: distraction with imagery versus general distraction. *Behav Res Ther.* 2002;40(3): 267–277. [https://doi.org/10.1016/S0005-7967\(01\)00012-2](https://doi.org/10.1016/S0005-7967(01)00012-2).
17. Morin C, Bootzin R, Buysse D, et al. Psychological and behavioral treatment of insomnia: update of the recent evidence. *Sleep.* 2006;29(11):1398–1414. <https://doi.org/10.1093/sleep/29.11.1398>.
18. Sivertsen B, Omvik S, Pallesen S, et al. Cognitive behavioral therapy vs zopiclone for treatment of chronic primary insomnia in older adults; a randomized controlled trial. *JAMA.* 2006;295(24):2851–2858. <https://doi.org/10.1001/jama.295.24.2851>.
19. Lastella M, Roach GD, Halson SL, Sargent C. Sleep/wake behaviours of elite athletes from individual and team sports. *Eur J Sport Sci.* 2015;15(2):94–100. <https://doi.org/10.1080/17461391.2014.932016>.
20. Hirshkowitz M, Whitton K, Albert SM, et al. National sleep foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health.* 2015;1(1):40–43.
21. Bruni O, Ottaviano S, Guidetti V, et al. The sleep disturbance scale for children (SDSC) construction and validation of an instrument to evaluate sleep disturbances in childhood and adolescence. *J Sleep Res.* 1996;5(4):251–261. <https://doi.org/10.1111/j.1365-2869.1996.00251.x>.
22. Tours JM, Muñíos R, Toro L. 9TH EUROPEAN CONGRESS OF PSYCHOLOGY (Granada 3-8 de julio de 2005). *Anuario de Psicología.* 2005;36(1):131–132 [fecha de Consulta 20 de Febrero de 2025]. ISSN: 0066-5126. Recuperado de: <https://www.redalyc.org/articulo.oa?id=97017363007>.
23. Merayo A, Gallego JM, Sans O, et al. Quantity and quality of sleep in young players of a professional football club. *Sci Med Footb.* 2022;6(4):539–544. <https://doi.org/10.1080/24733938.2021.1962541>.
24. Moseley L, Gradsar M. Evaluation of a school-based intervention for adolescent sleep problems. *Sleep.* 2009;32(3):334–341.
25. Milewski MD, Skaggs DL, Bishop GA, et al. Chronic lack of sleep is associated with increased sports injuries in adolescent athletes. *J Pediatr Orthop.* 2014;34(2): 129–133.

26. Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: a systematic literature review. *Sleep Med Rev.* 2015;21:50–58.
27. Bessot N, Moussay S, Sesboüé B, et al. The circadian rhythm of core temperature: effects of physical activity and aging. *Med Sci Sports Exerc.* 2006;38(6):1153–1160.
28. Vitale KC, Owens R, Hopkins SR, Malhotra A. Sleep hygiene for optimizing recovery in athletes: review and recommendations. *Int J Sports Med.* 2019;40(8):535–543. <https://doi.org/10.1055/a-0905-3103>. Epub 2019 Jul 9. PMID: 31288293; PMCID: PMC6988893.
29. Lacks P, Rotert M. Knowledge and practice of sleep hygiene techniques in insomniacs and good sleepers. *Behav Res Ther.* 1986;24(3):365–368.
30. Lastella A, Roach CL, Halson SM. The effect of age and sex on sleep quality in athletes. *J Sports Sci.* 2013;31(3):335–341. <https://doi.org/10.1080/02640414.2012.733822>.
31. Bolin DJ. Sleep deprivation and its contribution to mood and performance deterioration in college athletes. *Curr Sports Med Rep.* 2019;18(8):305–310. <https://doi.org/10.1249/JSR.0000000000000621>.
32. Driller MW, Lastella M, Sharp AP. Individualized sleep education improves subjective and objective sleep indices in elite cricket athletes: a pilot study. *J Sports Sci.* 2019;37(20):2121–2125. <https://doi.org/10.1080/02640414.2019.1620988>.
33. Nishinoue N, Takano T, Kaku A, et al. Effects of sleep hygiene education and behavioral therapy on sleep quality of white-collar workers: a randomized controlled trial. *Ind Health.* 2012;50(2):123–131.
34. Estivill-Domènech C, et al. Do elite athletes sleep well? *Apunts Medicina de l'Esport.* 2018;53(198):47–54.
35. Baron KG, Duffecy J, Berendsen MA, Cheung Mason I, Lattie EG. Feeling validated yet? A scoping review of the use of consumer-targeted wearable and mobile technology to measure and improve sleep. *Sleep Med Rev.* 2018;40:151–159.