



apunts

MEDICINA DE L'ESPORT

www.apunts.org



REVIEW

Importance of plantar sensitivity in the regulation of postural control and movement: Review

Caleb Araguas Garcia^{a,b,*}, Francisco Corbi Soler^b, Carles Vergés Salas^c

^a *Clinica podològica la Torrassa, Universitat de Barcelona, Barcelona, Spain*

^b *Departamento de Salud, Facultad de Ciencias de la Actividad Física y del Deporte, Centro de Lleida, Universitat de Lleida, Lleida, Spain*

^c *Ensenyament de Podologia, Departament de Ciències clíniques, Facultat de Medicina i Ciències de la Salut, Universitat de Barcelona, Barcelona, Spain*

Received 28 April 2016; accepted 12 September 2016

KEYWORDS

Sensitivity;
Skin;
Foot;
Sport;
Posture;
Injury

Abstract

The objective of this review is to analyse the influence of plantar sensitivity in improving the ability to re-balance and manage movement by analysing the mechanisms available to assess this and describing the factors that influence this. This project required a systematic search of the literature published between 2000 and 2016 in PubMed, Web of Science, and SPORTDiscus, as well as the references cited in relevant articles from these sources. The publication languages were Spanish and English, and a total of 9 items were collected and analysed.

Most studies suggest that sensory feedback from the foot is essential in maintaining general (postural and displacement) and specific (sport) patterns. Altering the quantity or quality of plantar afferent information not only appears to alter the creation of different patterns but may also increase the risk of injury.

© 2017 FC Barcelona. Published by Elsevier España, S.L.U. All rights reserved.

PALABRAS CLAVE

Sensibilidad;
Piel;
Pie;
Deporte;
Postura;
Lesión

Importancia de la sensibilidad plantar en la regulación del control postural y del movimiento: revisión

Resumen

El objetivo de esta revisión es analizar la influencia de la sensibilidad plantar en la mejora de la capacidad de reequilibrio y de gestión del movimiento, analizando los mecanismos que permiten valorarla y describiendo los factores que en ella influyen. Para ello se realizó una búsqueda sistemática de la bibliografía publicada entre 2000 y 2016, en las bases de datos PubMed, Web of Science, SPORTDiscus y en las referencias citadas

* Corresponding author.

E-mail address: calebaraguas@hotmail.com (C. Araguas Garcia).

en los artículos recuperados, cuyo idioma de publicación fuera español o inglés. Un total de 9 artículos fueron recuperados y analizados.

La mayoría de los estudios sugieren que la retroalimentación sensorial del pie resulta fundamental en el mantenimiento de los patrones generales (posturales y de desplazamiento) y específicos (deportivos). La alteración de la cantidad o calidad de la información aferente plantar no solo parece alterar la creación de los diferentes patrones, sino que podrá incrementar el riesgo de aparición de lesiones.

© 2017 FC Barcelona. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

The foot is a hypercomplex structure which supports the body and is characterised by being the only body part in contact with the ground. Among its various functions a key one is to provide afferent information to the central nervous system from plantar receptors, which will then be used to maintain posture and produce movement patterns. In recent years, several lines of research have focussed on the study of the influence that afferent information from the plantar receptors of the somatosensory system exerts on balance, postural control and movement, as well as on the occurrence of lesions.¹⁻⁴ Taking into account that the sole of the foot forms the point of contact between our body and the surface which supports it, and that it also has a large number of sensitive receptors, it is logical to assume that the sole of the foot will be directly involved in the management of posture and movement.^{5,6}

The skin is the largest organ in the human body.⁷ Its function is to act as a protective barrier, isolating the body from the surrounding environment, protecting it and contributing to maintaining its structures intact, while acting as a highly efficient communication system between the body and the immediate environment. In addition, the skin of the foot has certain characteristics that differentiate it from that of the rest of the human body: firstly, its epidermis presents greater thickness, which contributes to improving its resistance and its ability to cushion the body against loads. Secondly, the skin of the foot has no sebaceous glands, so that the hydrolipidic film is less rich in fatty components. Finally, this epidermis presents a large number of eccrine sweat glands, through which water, mineral salts and waste substances are excreted.⁸

The group of plantar receptors that provide the foot with sensitivity is heterogeneous due to the different types of skin receptors and the diversity of existing afferent nerve fibres.⁹ With the exception of nociceptors and thermoreceptors, most of the skin afferents correspond to the mechanoreceptors, which respond to pressure and vibration stimuli, and are related to myelinated afferent fibres (fast conducting fibres).¹⁰

Mechanoreceptors can be classified into two different groups: rapidly adapting and slowly adapting. Within the first group, we find Meissner's corpuscles and those of Paccini.¹¹ The former are also called tactile, and are located superficially (intervening in surface touch and responding to low-frequency vibrations).^{12,13} Paccini's corpuscles are found in the deep dermis of the skin and the interior of the adipose plantar tissue and respond quickly to sensations of deep pressure, high frequency vibration and stretching.¹⁴

In addition, we can find two types of slowly adapting mechanoreceptors: Merkel's disks and Ruffini's corpuscles.¹⁵ Merkel's disk is located on the surface of the epidermis of the skin and is involved in surface or discriminative touch. Ruffini's corpuscles are considered to be a variant of Meissner's corpuscles and are located more deeply, mediating sensations of thick and persistent touch.¹⁶

In addition, sensitivity levels on the plantar surface may vary greatly within the healthy¹⁷ and pathological population,¹⁸ with very different activation thresholds being observed depending on the age, area of the foot, sex or type of stimulus presented (vibratory or pressure-based).¹⁹ All this suggests that plantar afferent information could directly influence rebalancing ability and the creation of motor patterns.

Therefore, the main objective of this study is to analyse the influence of plantar sensitivity on changes in the ability to rebalance and manage movement, analysing the mechanisms that enable this to be assessed, and describing the factors that influence it.

Methodology

The different articles analysed in this study were located in the on-line computerised databases PubMed (Medline), Web of Science and SPORTdiscus. The keywords used for the search were: "sport injury", "sensitivity", "skin", "foot", and "posture". The search terms were initially used independently and were subsequently grouped using the terms *and* and *or*. Regarding the selection criteria, we included experimental and quasi-experimental studies in English, published in impact journals. The search for articles was restricted to the period from January 2000 to January 2016. In addition, regarding the remaining inclusion criteria, the PICR (Participants, Intervention, Comparison, Results) reference was followed. The articles included were organised using "Mendeley" publication management software. Once the different studies had been analysed, duplicates were discarded; studies with little methodological rigour; case studies; or those where the subject matter did not exactly fit our study objectives; and those whose text could not finally be obtained.

Results

The article search and selection strategy adopted in this review obtained a total of 9 articles which finally met the inclusion criteria. The characteristics of the different studies identified can be found in Table 1.

Several different methodologies have been used for the assessment of plantar sensitivity: tactile,²⁹ algescic,³⁰ thermal,³¹ vibratory,³² baresthetic,³³ proprioception,³⁴ reflex³⁵ (Table 2). In general, in these methodologies, mechanical stimuli of different intensity are usually applied at symmetrical points on both hemispheres of the body, or the level of sensitivity between different body areas is compared.³⁶ To achieve this, it appears to be important that the assessments are performed in the absence of fatigue, since it has been evidenced that the existence of fatigue seems to have a negative influence on assessment.³⁷ Although many of these methodologies come from the clinical field,³⁸ many can be applied directly or with slight modifications in healthy subjects and sportsmen.³⁹

External factors influencing plantar sensitivity

Temperature

Several studies have suggested that the local application of cold seems to acutely increase sympathetic nervous activity in the applied area;⁴⁰ to produce local reflex vasoconstriction;⁴¹ to reduce the skin temperature,⁴² and to alter local sensitivity.⁴³ On this point, Nurse and Nigg⁶ found that the pressure centre moved from areas where cold was applied to areas where sensitivity was greater, and variations in patterns of muscle activity were recorded. On the other hand, Eils et al.,⁴⁴ following a similar methodology, found reductions in the peaks of plantar pressure on stimulated areas and an alteration in plantar pressure patterns. This fact can be interpreted as a form of body protection, where the body attempts to avoid contact in areas of the foot where sensitivity is altered, seeking a support that searches for areas with greater sensitivity.

Footwear

Several authors have theorised about the influence that sport footwear could have on changes in plantar sensitivity as regards the information provided afferently from the plantar receptors⁴⁵ and the reduction of their stimulation threshold.⁴⁶ Factors such as hardness, design or thickness of the sole of the footwear,⁴⁷ the type of footwear,⁴⁸ the type of insole,⁴⁹ and the use of certain specific footwear, such as football boots,⁵⁰ appear to directly influence the level of plantar sensitivity. In addition, it has been shown that the use of air pocket footwear could increase the instability level in the foot and increase the risk of injury, as a result of the reduction in afferent information.⁵¹ It has also been seen that certain footwear, such as minimalist footwear, could also have repercussions on plantar sensitivity levels,⁵² especially when this provokes painful stimuli.⁵³ On this point, it is clear that the type and characteristics of footwear used could produce changes in the quantity and quality of the information collected.⁵⁴

Type of contact surface

The hardness of the terrain or footwear are factors that can affect plantar sensitivity and influence the pattern of

movement.⁵⁵ Chiang and Wu⁵⁶ observed that as soil hardness reduced to a softer surface, body stability decreased, afferent response times increased, and plantar pressures recorded changed, all of which resulted from changes in skin receptors rather than muscle receptors.⁵⁷

Hyperkeratosis

Hyperkeratosis occurs when there is thickening of the corneal layer of the epidermis caused by hypertrophy (enlargement of its cells) or hyperplasia (increase in the number of cells). This increase in the size or number of cells mainly affects the keratinocytes or corneocytes, which are the most numerous cells in the outermost layer of the epidermis.⁵⁸ Plantar hyperkeratosis is able to inhibit sensory feedback in the area where it develops,⁵⁹ which will cause displacement of the centre of pressures from the areas with less sensitivity to those with greater sensitivity.⁶⁰

Mechanical vibrations

Although the vast majority of studies on the vibrotactile perception of the skin have been run on the hand, there is a high degree of consensus as to the influence that mechanical vibrations have on the foot, especially in certain pathologies such as the diabetic foot. This is due to the large number of mechanoreceptors present in the sole of the foot (Kennedy and Inglis⁶¹ located 104 mechanoreceptors in the skin of the human foot), and to the existence of specific vibration thresholds depending on the area of the foot (lower vibrotactile thresholds have been found in the midfoot than in the heel or toes).¹⁷ All this will affect the existence of specific efferent responses, depending on the type of vibratory stimulus presented. In this direction, the use of mechanical vibrations as a training method has been found to alter the level of plantar sensitivity and affect rebalancing ability,⁶² especially in the 10 min after its application and when mechanical vibrations are high frequency.⁶³

Proprioceptive insoles

It has recently been suggested that the placement of insoles of different textures seems to improve postural control,⁶⁴ increasing the ankle's ability for positional discrimination, preventing the appearance of injuries⁶⁵ and creating more efficient muscle patterns.⁶⁶ In addition, these improvements seem to appear in a relatively short time (5 weeks).⁶⁴ Santos et al.⁶⁷ found that the use of football boots decreased plantar sensitivity due to a reduction in the foot-ground contact area and an increase in plantar pressure peaks, and that when football boots were used in combination with stimulatory insoles, sensitivity and proprioceptive response improved considerably.

Discussion

Traditionally, the foot has been considered as a fundamental link within the maintenance of postural balance.⁶⁸ The first models that attempted to explain how human beings

Table 1 Characteristics of the studies run on changes in plantar sensitivity in the foot

Author (year)	Number sample	Age (years)	Objective	Instrument assessment	Mechanism of change in sensitivity	Area validated	Results
Kavounoudias and Roll (1998) ²⁰	10	Not specified	To study the effect of the application of vibrations in the sole of the foot on body balance	Oscillometers	Vibration	Foot	The stimulus caused the body to lean over sideways involuntarily, always in the opposite direction to the area of the foot where the vibration was applied
Nurse and Nigg (1999) ²¹	15	Not specified	To analyse the relationship between sensitivity (to pressure and vibration) and pressures recorded in the sole of the foot during the gait	Semmes-Weinstein monofilaments, oscillometer	Pressure vibration	Hallux	There is an inverse correlation between the vibration threshold of the hallux at high frequencies and the maximum peak of plantar pressure detected in the hallux during gait and sprint. Subjects with greater sensitivity in the hallux presented higher peaks of plantar pressure during race
Bensmaïa et al. (2005) ²²	22	Not specified	To investigate the effects on the sensitivity of suprathreshold vibratory stimulation	Linear motor of controlled feedback	Vibration	Foot	The application of vibratory stimuli above the stimulation threshold can alter the activation capacity of the afferent mechanoreceptors
Eils et al. (2002) ²³	40	25,3 ± 3,3	To determine the influence of the reduction of plantar sensitivity by reducing foot temperature in the distribution of plantar pressure patterns during gait	Semmes-Weinstein monofilaments	Pressure	Sole of foot	The reduction in skin temperature modifies the patterns of distribution in the pressure of the sole of the foot along its support
Wang and Lin (2007) ²⁴	21	Not specified	To investigate the correlation between the degree of somatosensory loss in the sole of the foot and the degree of postural stability	Not specified	Tactile	Sole of foot	Ischemia is associated with an increase in the loss of cutaneous plantar sensitivity and an increase in postural instability

Table 1 (continue)

Author (year)	Number sample	Age (years)	Objective	Instrument assessment	Mechanism of change in sensitivity	Area validated	Results
Meyer et al. (2004) ²⁵	6	26 ± 10	To associate the influence of local injected subcutaneous anaesthesia on skin plantar sensation and body balance	Not specified	Tactile	Sole of the foot	Local anesthesia is associated with increased skin plantar sensitivity loss and an increase in postural instability, mainly when no information can be provided by the eye
Burcal et al. (2016) ²⁶	45	20,2 ± 2,8	To determine if there are any differences in plantar sensation between runners who have suffered ankle injuries and those who have not	Electric instrument with sensorial threshold	Vibration	Plantar arch	Runners with a history of injury showed an increase in the vibration threshold in the region of the arch when compared with uninjured runners. A higher plantar sensitivity alters sensory functions after the injury. This factor can influence underlying postural control and contribute to altered load responses in injured runners
Vie et al. (2014) ²⁷	30	28,7 ± 8,6	To analyse whether somatosensory plantar sensation could be affected by the use of hard metatarsal pads	Stevens Power Function	Pressure	Forefoot	Metatarsal pads with a relatively high level of hardness increased the perception of mechanical stimulation in the forefoot, as compared with soft metatarsal pads
Lipsitz et al. (2014) ²⁸	12	62,3 ± 4,5	To ascertain whether sub sensory vibrational noise applied to the sole of the foot using a piezoelectric vibratory insole could improve sensitivity and balance	Urethane foam insole with piezoelectric sensor	Vibration	Sole of the foot	The application of the principle of stochastic resonance for the sensory system of the foot, using a new low tension piezoelectric technology can improve measurements of balance and gait

Table 2 Methodologies for assessment of plantar sensitivity

Author and year	Name	Instrument to be used	Protocol	Warning signs
Pedro et al. (2011) ²⁹	Tactile epicritical sensitivity	Fine brush cotton wool	With the subject lying on an examination table, and deprived of vision. Using a fine brush or cotton wool, different areas of the feet will be brushed lightly, the subject having to identify when they are touched	The subject is not able to recognize any sensation at different points of the foot
Runkle et al. (2006) ³⁰	Algesic sensitivity	Blunt-ended pin	Subject lying on an exploratory stretcher and deprived of vision. Using a pin with a blunt tip, different areas of the foot are pressed and the patient is asked if they feel pain	If the subject does not detect stimulation at more than one point on each foot
Cacicedo et al. (2011) ³¹	Thermal sensitivity	Thermal bar	Apply the thermal bar on different areas of the foot, always avoiding injured areas of those with hyperkeratosis	When the temperature applied to the foot of the subject is higher than 40 °C or below 25 °C and the subject is unable to identify this
Fernández et al. (2009) ³²	Vibratory sensitivity	Rydel Seiffer graduated tuning fork (128 Hz)	With the subject lying supine on an examination table, a tuning fork will be made to vibrate and placed on the tip of the first toe at the head of the first metatarsal and will be moved in the direction of the tibial and peroneal malleoli	The moment the subject indicates that he/she has ceased to feel the vibration, the reading that appears at the point of intersection of the two triangles will be noted
Gutiérrez et al. (2005) ³³	Barethetic or pressor sensitivity	Semmens-Weinstein 5.07 monofilament	The subject is placed in supine position on an examination table. The nurse will press perpendicularly with the monofilament on the skin without sliding over it. The thread should be folded for 1 to 1.5 seconds	When the patient is not able to feel one of the points where the monofilament is bent, this means that the tactile sensation has decreased below the threshold of sensitive protection
Sullivan et al. (2001) ³⁴	Proprioception	-	A joint or limb will be placed in a given position and the participant will be asked to reproduce the same position in the contralateral limb	Not being able to reproduce the position with the contralateral limb
Singh et al. (2005) ³⁵	Achilles reflex	Neurological percussion hammer	A small blow is struck with the neuropercutor hammer on the Achilles tendon to provoke a flexor plantar response of the foot	The absence of reflex response indicates that the patient is suffering from neuropathic involvement

rebalanced presented the body as an inverted pendulum⁶⁹ whose axis of rotation was located in the ankle and where the constant wavering of the centre of gravity was readjusted, thanks to the information provided through afferent pathways.⁷⁰ Although other explanatory models were later

presented that attributed a fundamental role to the hip in the rebalancing process,⁷¹ at present there is a certain degree of consensus as to the existence of a combined and intermittent model where there is specific interaction between different body parts involved in a posture or move-

ment.⁷² This will be configured, to a large extent, from the information provided by the somatosensory system.⁷³

On this point it seems that the information provided by the plantar receptors is fundamental, since it seems that any alteration in the sensitivity of the skin receptors in the sole of the foot could have direct involvement in the alteration of the kinetic and muscular support patterns.⁷⁴ For example, several studies have observed that, in subjects subjected to weightlessness, the loss of afferent information provided by the sole of the foot could lead to a decrease in the activity of the tonic muscles.⁷⁵ Kozlovskaya et al.⁷⁶ found that by submerging a group of subjects inside a tank of water for an extended period of time, there was a decrease in the activation levels of the extensional postural tonic muscles⁷⁷ and an increase in those of the flexor,⁷⁸ as a consequence of the removal of the afferent information provided by the sole of the foot. Although many of these changes are going to occur locally, all this suggests that changes could also be produced at the brain level.⁷⁹ For example, Liepert et al.⁸⁰ found that after 4-6 weeks of immobilisation of the foot as a consequence of a fracture, there was a decrease in cortical representation in the brain in the area corresponding to the foot.

Similarly, it appears that a decrease in sensory feedback levels in selective areas of the foot during gait could cause displacement of the centre of plantar pressure from areas where there is a loss of sensitivity to areas of the sole of the foot with greater sensitivity.⁸¹ This may produce changes in plantar pressure patterns (peak pressure and pressure-time integral). Moreover, it seems that all this could have repercussions on the muscles involved in the movement and their level of activity.⁸²

On the contrary, several studies suggest that certain activities may favour the local ability to collect afferent information or the central ability to activate specific brain areas, generating specific adaptations resulting from training. Meier et al.⁷⁹ found that, when comparing professional handball players with professional dancers, the latter were able to optimise the level of activation of specific areas of the brain related to segmentary control of the foot. This could be due, at least in part, to the development of a greater sensory ability developed in the dancers, due to working barefoot, adopting different positions on the foot in situations of instability and stimulating very reduced areas of contact. Similarly, there have also been improvements in levels of brain activation in athletes who use their feet specifically in their sports. In a study comparing the football player, Neymar Jr., with other football players (three professionals and one amateur) and two professional swimmers, it was observed that Neymar had less neural activity in the area of his brain related to the foot when making simple movements with it, which was interpreted as a sign of efficiency.⁸³ In addition, these results are in line with those obtained in other bodily regions such as hands, in groups that usually have to develop high levels of performance, such as pianists or keyboard players.⁸⁴ This has also been observed in primates.⁸⁵

All of this suggests that any alteration of the information provided afferently from the feet could modify the patterns of muscle activation during standing,⁴¹ directly influencing postural stability levels.

On the other hand, several studies have shown the existence of a direct relationship between a decrease in plantar sensitivity and increased risk of lesion. For example, it has been observed that sportsmen who have suffered ankle injuries have lower levels of postural stability,⁸⁶ a decrease in joint stability levels, and altered motor patterns.⁸⁷ Similarly, Steinberg et al.⁸⁸ observed that in patients with anterior cruciate ligament injury there was poorer vibratory sensitivity in the foot and ankle.

These changes appear to be the result of the appearance of a change in the functioning of the afferent information channels, provoked by the lesion, which justifies the need to re-educate and optimise its functioning. On the other hand, the possible negative influence that cryotherapy (as a treatment or recovery method) may have on proprioceptive ability has been questioned, verifying that the effects on proprioceptive ability may be variable, depending on the time of application⁸⁹ and the body area analysed.⁹⁰

Several authors have researched the effect of proprioceptive ankle training on static body balance, noting that in healthy individuals, this produces positive effects.⁹¹ Han et al.⁹² measured the ankle's proprioceptive ability in 100 elite athletes from 5 different sports (gymnastics, football, swimming, badminton and dance), observing a direct relationship between the level of ankle proprioception and performance in sport, even at Olympic level. All this appears to indicate how obtaining improvements in body balance through plantar sensitivity enables sportsmen to reduce the level of attention they have to pay, consciously or unconsciously, to the maintenance of stability, thus being able to devote more attention to improvement in their technique. In a study comparing gymnasts with sportsmen who did not specifically work on proprioception, it was observed that the former decreased their dependence on postural control processes when they were subjected to different tasks or tasks of greater complexity.⁹³ Therefore, it seems logical to think that specific training of plantar receptors could have positive effects on creating and optimising motor patterns. Improvement in this regard should therefore be considered in both sport and preventive and injury treatment programmes.

Conclusions and recommendations

As suggested by our review, sensory feedback from the foot is essential in the maintenance of general (postural and displacement) and specific (sport) patterns. Changes in the quantity or quality of plantar afferent information will not only upset the creation of different patterns but may also increase the risk of injury. In this sense, it will be fundamental to maintain and optimise the ability to collect afferent information from the different systems, as a way to avoid the appearance of injuries and pathologies and improve performance. Continuous monitoring of information gathering levels, coupled with the creation of stimulation tasks, are alternatives that should be considered.

Conflict of interest

The authors declare no conflict of interest.

References

- Billot M, Handrigan GA, Simoneau M, Teasdale N. Reduced plantar sole sensitivity induces balance control modifications to compensate ankle tendon vibration and vision deprivation. *J Electromyogr Kinesiol.* 2015;25:150-60.
- Powell MR, Powden CJ, Houston MN, Hoch MC. Plantar cutaneous sensitivity and balance in individuals with and without chronic ankle instability. *Clin J Sport Med.* 2014;24:490-6.
- Fort A, Romero D. Rol del sistema sensoriomotor en la estabilidad articular durante las actividades deportivas. *Apunts Med Esport.* 2013;48:69-71.
- Robbins SE, Hanna AM, Govw G. Overload protection avoidance response to heavy plantar surface loading. *Med Sci Sports Exerc.* 1988;20:85-92.
- Corbi F. Análisis de las presiones plantares y su relación con la velocidad de la pelota durante el goleo paralelo de derecha en tenis [tesis doctoral]. Barcelona: Universidad de Barcelona; 2008.
- Nurse MA, Nigg BM. The effect of changes in foot sensation on plantar pressure and muscle activity. *Clin Biomech (Bristol).* 2001;16:719-27.
- Freedberg A, Fitzpatrick's I. *Dermatology in General Medicine.* 5th ed. New York: McGraw-Hill; 1999.
- Fustero I. Cuidado de los pies. *Offarm.* 2007;2:66-72.
- Dupui PH, Costes-Salon MC, Montoya R, Séverac A, Lazerges M, Pagès B, et al. Intérêt de l'analyse fréquentielle des oscillations posturales lors de l'équilibre dynamique. Société d'études et de recherches multidisciplinaires sur la locomotion. Paris: Collège de France. 1990.
- Dupui PH, Montoya R. Approche physiologique des analyses posturographiques statiques et dynamiques. *Physiologie, Techniques, Pathologies.* 3me ed. Paris: Posture et Equilibre; 2003.
- Vega JA, García-Suárez O, Montaña JA, Pardo B, Cobo JM. The Meissner and Pacinian sensory corpuscles revisited new data from the last decade. *Microsc Res Tech.* 2009;72:299-309.
- Hoffmann JN, Montag AG, Dominy NJ. Meissner corpuscles and somatosensory acuity: The prehensile appendages of primates and elephants. *Anat Rec A Discov Mol Cell Evol Biol.* 2004;281:1138-47.
- Luo W, Enomoto H, Rice FL, Milbrandt J, Ginty DD. Molecular identification of rapidly adapting mechanoreceptors and their developmental dependence on ret signaling. *Neuron.* 2009;64:841-56.
- Rein S, Manthey S, Zwipp H, Witt A. Distribution of sensory nerve endings around the human sinus tarsi: A cadaver study. *J Anat.* 2014;224:499-508.
- Maeda T, Ochi K, Nakakura-Ohshima K, Youn SH, Wakisaka S. The Ruffini ending as the primary mechanoreceptor in the periodontal ligament: Its morphology, cytochemical features, regeneration, and development. *Crit Rev Oral Biol Med.* 1999;10:307-27.
- Thibodeau GA, Patton KT. *Anatomía y fisiología.* 6th ed. Madrid: Elsevier; 2007.
- Kekoni J, Hämmäläinen H, Rautio J, Tukeya T. Mechanical sensitivity of the sole of the foot determined with vibratory of varying frequency. Experimental brain research. *Experimentelle Hirnforschung. Exp Brain Res.* 1989;78:419-24.
- Rautio J, Kekoni J, Hämmäläinen H, Härmä A. Mechanical sensitivity in free and island flaps of the foot. *J Reconstr Microsurg.* 1989;5:119-25.
- Hennig EM, Sterzing T. Sensitivity mapping of the human foot: Thresholds at 30 skin locations. *Foot Ankle Int.* 2009;30:986-91.
- Kavounoudias A, Roll R, Roll JP. The plantar sole is a 'dynamic map' for human balance control. *Neuroreport.* 1998;9:3247-52.
- Nurse MA, Nigg BM. Quantifying a relationship between tactile and vibration sensitivity of human foot with plantar pressure distributions during gait. *Clin Biomech (Bristol).* 1999;14:667-72.
- Bensmaïa SJ, Leung YY, Hsiao SS, Johnson KO. Vibratory adaptation of cutaneous mechanoreceptive afferent. *J Neurophysiol.* 2005;94:3023-36.
- Eils E, Nolte S, Tewes M, Thorwesten L, Olker KV, Rosenbaum D. Modified pressure distribution patterns in walking following reduction of plantar sensation. *J Biomech.* 2002;35:1307-13.
- Wang TY, Lin S. Sensitivity of plantar cutaneous sensation and postural stability. *Clin Biomech.* 2008;23:493-9.
- Meyer F, Oddsson IE, de Luca J. Reduced plantar sensitivity alters postural responses to lateral perturbations of balance. *Exp Brain Res.* 2004;157:526-36.
- Burcal CJ, Wikstrom EA. Plantar cutaneous sensitivity with and without cognitive loading in people with chronic ankle instability, copers, and uninjured controls. *J Orthop Sports Phys Ther.* 2016;1-24.
- Vie B, Loffredo R, Sanahdji F, Weber JP, Jammes Y. Consequences of repetitive toenail cutting by podiatric physicians on force production, endurance to fatigue, and the electromyogram of the flexor digitorum superficialis muscles. *J Am Podiatr Med Assoc.* 2014;104:486-92.
- Lipsitz LA, Lough M, Niemi J, Trivison T, Howlett H, Manor B. A shoe insole delivering subsensory vibratory noise improves balance and gait in healthy elderly people. *Arch Phys Med Rehabil.* 2015;96:432-9.
- Pedro J, Rivera AM, Olivan RM. Manejo del pie diabético. *Guía de Enfermería.* 2011;79:29-41.
- Runkle I, Díaz J, Durán A, Romero L. Técnicas de exploración de la sensibilidad en la patología del pie. *Av Diabetol.* 2006;22:42-9.
- Cacicedo R, Castañeda C, Cosío F, Delgado A, Fernández B, Gómez M, et al. Manual de prevención y cuidados locales de heridas crónicas. Cantabria: Servicio Cántabro de Salud; 2011.
- Fernández T, Montequín F. Manual para la prevención, diagnóstico y tratamiento del pie diabético. *Rev Cuba Angiol Cirug Vasc.* 2009;10:42-96.
- Gutiérrez T, Reyes R, Pizarro C. Pie diabético, una visión fisiátrica. *Rev Hosp Clin Univ Chile.* 2005;16:14-23.
- O'Sullivan S. *Assessment of Motor Function.* Philadelphia: Physical Rehabilitation; 2005.
- Singh N, Armstrong D, Lipsky B. Preventing foot ulcers in patients with diabetes. *JAMA.* 2005;293:217-28.
- Bates B, Bickley LS, Hoekelman RA. *A guide to physical examination and history taking.* 6th ed. Philadelphia: JB Lippincott; 1995.
- Sweeny VP. History and physical examination in patients with neurologic complaints. In: Kelley WN, De Vita VT, editors. *Textbook of Internal Medicine.* 2nd ed. Philadelphia: JB Lippincott; 1992. pp. 2251-5.
- Murray J. Neurologic history and examination. In: Stein JH, editor. *Internal Medicine.* 4th ed. St Louis: Mosby-Year Book; 1994. pp. 960-1.
- Martin JB, Hauser SL. Estudio del paciente con una enfermedad neurológica. In: Braunwald E, editor. *Harrison, principios de medicina interna.* 15th ed. México DF: McGraw-Hill Interamericana; 2002. pp. 2219-25.
- Ishida K, Nakamura T, Kimura K, Kanno N, Takahashi N, Kamijo Y-I, et al. Suppression of activation of muscle sympathetic nerve during non-noxious local cooling after the end of local cooling in normal adults. *Eur J Appl Physiol.* 2016;116:851-8.
- Diener J, Dichgans B, Guschlbauer H. The significance of proprioception on postural stabilization as assessed by ischemia. *Brain Res.* 1984;296:103-9.
- Zaproudina N, Lipponen JA, Eskelinen P, Tarvainen MP, Karjalainen PA, Närhi M. Measurements of skin temperature re-

- sponses to cold exposure of foot and face in healthy individuals: Variability and influencing factors. *Clin Physiol Funct Imaging*. 2011;31:307-14.
43. Eglin CM, Golden FS, Tipton MJ. Cold sensitivity test for individuals with non-freezing cold injury: The effect of prior exercise. *Extrem Physiol Med*. 2013;2:16.
 44. Eils E, Nolte S, Tewes M, Thorwesten L, Olker KV, Rosenbauma D. Modified pressure distribution patterns in walking following reduction of plantar sensation. *J Biomech*. 2002;35:1307-13.
 45. Robbins SE, Hanna AM. Running-related injury prevention through barefoot adaptations. *Med Sci Sports Exerc*. 1987;19:148-52.
 46. Robbins SE, Hanna AM, Jones L. Sensory attenuation induced by modern athletic footwear. *J Test Eval*. 1988;16:422-6.
 47. Losa ME, Becerro R, Palacios D. Impact of soft and hard insole density on postural stability in older adults. *Geriatr Nurs*. 2012;33:264-71.
 48. Schlee G, Sterzing T, Milani TL. Effects of footwear on plantar foot sensitivity: A study with Formula 1 shoes. *Eur J Appl Physiol*. 2009;106:305-9.
 49. Qu X. Impacts of different types of insoles on postural stability in older adults. *Appl Ergon*. 2015;46 Pt A:38-43.
 50. Waddington G, Adams R. Football boot insoles and sensitivity to extent of ankle inversion movement. *Br J Sports Med*. 2003;37:170-5.
 51. Serrano MA. Analysis of the instability of the foot with sport footwear or air chamber, during statics. *Rev Int Pod*. 2007;1:7-15.
 52. Trombini-Souza F, Matias AB, Yokota M, Butugan MK, Goldenstein-Schainberg C, Fuller R, et al. Long-term use of minimal footwear on pain, self-reported function, analgesic intake, and joint loading in elderly women with knee osteoarthritis: A randomized controlled trial. *Clin Biomech*. 2015;30:1194-201.
 53. Ryan M, Elashi M, Newsham-West R, Taunton J. Examining injury risk and pain perception in runners using minimalist footwear. *Br J Sports Med*. 2014;48:1257-62.
 54. Khoury M, Wolf A, Debbi EM, Herman A, Haim A. Foot center of pressure trajectory alteration by biomechanical manipulation of shoe design. *Foot Ankle*. 2013;34:593-8.
 55. Chambon N, Delattre N, Guéguen N, Berton E, Rao G. Is midsole thickness a key parameter for the running pattern? *Gait Posture*. 2014;40:58-63.
 56. Chiang J, Wu G. The effects of surface compliance on foot pressure in stance. *Gait Posture*. 1996;4:122-9.
 57. Chiang J, Wu G. The influence of foam surfaces on biomechanical variables contributing to postural control. *Gait Posture*. 1997;5:239-45.
 58. Rubin L. Hyperkeratosis in response to mechanical irritation. *J Invest Dermatol*. 1949;13:313-5.
 59. Balanowski KR, Flynn LM. Effect of painful keratoses debridement on foot pain, balance and function in older adults. *Gait Posture*. 2005;22:302-7.
 60. Kavounoudias A, Roll R, Roll J. The plantar sole is a dynamometric map for human balance control. *Neuroreport*. 1998;8:3247-52.
 61. Kennedy PM, Inglis JT. Interaction effects of galvanic vestibular stimulation and head position on the soleus H reflex in humans. *Clin Neurophysiol*. 2002;113:1709-14.
 62. Sonza A, Maurer C, Achaval M, Zaro MA, Nigg BM. Human cutaneous sensors on the sole of the foot: Altered sensitivity and recovery time after whole body vibration. *Neurosci Lett*. 2013;533:81-5.
 63. Sonza A, Robinson CC, Achaval M, Zaro MA. Whole body vibration at different exposure frequencies: Infrared thermography and physiological effects. *ScientificWorldJournal*. 2015;2015:452657.
 64. Steinberg N, Waddington G, Adams R, Karin J, Begg R, Tirosh O. Can textured insoles improve ankle proprioception and performance in dancers? *J Sports Sci*. 2015;30:1-8.
 65. Steinberg N, Waddington G, Adams R, Karin J, Tirosh O. The effect of textured ballet shoe insoles on ankle proprioception in dancers. *Phys Ther Sport*. 2016;17:38-44.
 66. Kelleher KJ, Spence WD, Solomonidis S, Apatsidis D. The effect of textured insoles on gait patterns of people with multiple sclerosis. *Gait Posture*. 2010;32:67-71.
 67. Santos D, Carline T, Flynn L, Pitman D, Feeney D, Patterson C, et al. Distribution of in-shoe dynamic plantar foot pressures in professional football players. *Foot*. 2001;11:10-4.
 68. Watanabe I, Okubo LJ. The role of plantar mechanoreceptor in the equilibrium control. In: Colen B, editor. *Vestibular and Oculomotor Physiology: International Meeting of the Baray Society*. New York: Annals of the New York Academy of Sciences; 1981. pp. 855-64.
 69. Kuo AD. The six determinants of gait and the inverted pendulum analogy: A dynamic walking perspective. *Human Movement Science*. 2007;26:617-56.
 70. Roberts A, Stenhouse E. The nature of postural sway. *Agressologie*. 1976;17A:11-4.
 71. Horak FB, Nashner LM. Central programming of postural movements: adaptation to altered support surface configurations. *J Neurophysiol*. 1986;55:1369-81.
 72. Kilby MC, Molenaar PCM, Newell KM. Models of postural control: Shared variance in joint and COM motions. *PLoS One*. 2015;10:e0126379.
 73. Manchester D, Woollacott M, Zederbauer-Hylton N, Marin O. Visual, vestibular and somatosensory contributions to balance control in the older adult. *J Gerontol*. 1989;44:M118-27.
 74. Chen H, Nigg B, Hulliger M, de Koning J. Influence of sensory input on plantar pressure distribution. *Clin Biomech*. 1995;10:271-4.
 75. Grigoriev AI, Kozlovskaya IB, Shenkman BS. The role of support afferents in organization of the tonic muscle system. *Russ J Physiol*. 2004;90:37-43.
 76. Kozlovskaya IB, Tomilovskaya ES, Berger M, Gerstenbrand F. Alterations of characteristics of horizontal gaze fixation reaction in long-term space flights. *J Gravit Physiol*. 2007;14:P79-80.
 77. Koryak YA. Electromyographic study of the contractile and electrical properties of human triceps surae muscle in a simulated microgravity environment. *J Physiol*. 1998;510:287-95.
 78. Miller TF, Saenko IV, Popov DV, Vinogradova OL, Kozlovskaya IB. Effect of mechanical stimulation of the support zones of soles on the muscle stiffness in 7-day dry immersion. *J Gravit Physiol*. 2003;10:61-2.
 79. Meier J, Topka MS, Hänggi J, Meier J, Topka MS. Differences in cortical representation and structural connectivity of hands and feet between professional handball players and ballet dancers. *Neural Plast*. 2016;2016:1-17.
 80. Liepert J, Tegenthoff M, Malin J-P. Changes of cortical motor area size during immobilization. *Electroencephalogr Clin Neurophysiol Mot Control*. 1995;97:382-6.
 81. Kavounoudias A, Roll R. The plantar sole is a dynamometric map for human balance control. *Neuroreport*. 1998;9:3247-52.
 82. Magnusson M, Enbom H, Johansson R, Pyykkö I. Significance of pressor input from the human feet in anterior-posterior postural control. *Acta Otolaryngol*. 1990;110:182-8.
 83. Naito E, Hirose S. Efficient foot motor control by Neymar's brain. *Front Hum Neurosci*. 2014;8:594.
 84. Jäncke L, Shah NJ, Peters M. Cortical activations in primary and secondary motor areas for complex bimanual movements in professional pianists. *Brain Res Cogn Brain Res*. 2000;10:177-83.
 85. Picard N, Matsuzaka Y, Strick PL. Extended practice of a motor skill is associated with reduced metabolic activity in M1. *Nat Neurosci*. 2013;16:1340-7.
 86. Lin CF, Lee IJ, Liao JH, Wu HW, Su FC. Comparison of postural stability between injured and uninjured ballet dancers. *Am J Sports Med*. 2011;39:1324-31.

87. Lin CW, Su FC, Lin CF. Influence of ankle injury on muscle activation and postural control during ballet grand plié. *J Appl Biomech.* 2014;30:37-49.
88. Steinberg N, Waddington G, Adams R, Karin J, Tirosh O. Use of a textured insole to improve the association between postural balance and ankle discrimination in young male and female dancers. *Med Probl Perform Art.* 2015;30:217-23.
89. Thain PK, Bleakley CM, Mitchell AC. Muscle reaction time during a simulated lateral ankle sprain after wet-ice application or cold-water immersion. *J Athl Train.* 2015;50:697-703.
90. Uchio Y, Ochi M, Fujihara A, Adachi N, Iwasa J, Sakai Y. Cryotherapy influences joint laxity and position sense of the healthy knee joint. *Arch Phys Med Rehabil.* 2003;84:131-5.
91. Karakaya MG, Rutbl H, Akpınar E, Yildirim A, Karakaya IC. Effect of ankle proprioceptive training on static body balance. *J Phys Ther Sci.* 2015;27:3299-302.
92. Han J, Anson J, Waddington G, Adams R. Sport attainment and proprioception. *Int J Sports Sci Coach.* 2014;9:159-70.
93. Davlin CD. Dynamic balance in high level athletes. *Mot Skills.* 2004;98:1171-6.