



REVIEW

Intractable lateral epicondylitis: A differential diagnosis algorithm for a correct clinical interpretation



Jose Miguel Aguililla Liñan^{a,*}, Maria Isabel Miguel Pérez^b, Jordi Palau González^c, Ingrid Möller Parera^b

^a Gimbernat School of Physiotherapy, Universitat Autònoma de Barcelona, Sant Cugat del Vallès 08174, Spain

^b Department of Pathology and Experimental Therapeutics, Human Anatomy and Embryology Unit, Faculty of Medicine and Health Sciences, University of Barcelona, Barcelona, Spain

^c Centre for Hand Therapy (Centre de la Mà de Barcelona), Barcelona, Spain

Received 12 November 2020; accepted 22 February 2021

KEYWORDS

Tennis elbow;
Chronic pain;
Diagnosis;
Physical examination

Abstract The epicondylalgia is the most frequent upper extremity pathology in adults and it can become an “intractable lateral epicondylitis” when patients do not improve with the treatment received. This is a complex entity that includes several musculo-tendinous, articular and neural syndromes than can coexist and they can also be confused with each other. For this reason, it is necessary to do a systematized and exhaustive evaluation where all the dysfunctions capable of generating the symptoms are precisely and independently analyzed. On this basis, a 7 steps assessment algorithm is proposed on this paper to enable the clinician to perform a complete and organized evaluation of these patients, to achieve a correct clinical interpretation.

Background

Lateral elbow pain is a prevalent and highly unspecific finding that can have different causes. In spite of being a common condition, its chronification rate is high¹, and it is known as *intractable lateral epicondylitis* when the patient does not seem to get better with treatment². It is often believed that, in these patients, their symptoms result from epicondylitis, commonly known as ‘tennis elbow’^{3–6}. Nevertheless, lateral elbow pain is a heterogeneous entity in

its clinical presentation and its pathophysiology where the tendinous pathology can also involve changes in the processing of nociceptive information and impairments in motor and sensory function⁷. Apart from this, this tendinous pathology can also coexist with and/or be mistaken for other clinical syndromes of the elbow^{8,9} of neural^{10,11}, articular^{12–14}, and/or muscular^{15,16} origin, in which psychosocial factors can play a key role^{7,17}. All these aspects make intractable lateral epicondylitis a challenging condition for clinicians since there is no consensus on its diagnosis³.

This paper aims to be a guideline for all those healthcare providers who have to face the challenge of diagnosing these patients. The reader will find a seven-step algorithm through which they will apply their clinical reasoning and decision

* Corresponding author.

E-mail address: jose.aguililla@eug.es (J.M. Aguililla Liñan).

making skills when assessing the patient. The clinician will disclose the different causes of epicondylalgia in an orderly and clear way, starting with the most frequent causes and moving to the least frequent ones, allowing them to make a better clinical interpretation.

First step: confirming or excluding tendinopathy

Palpating the different anatomical structures in the lateral region of the elbow when carrying out a physical examination is very important for the differential diagnosis of *intractable lateral epicondylitis*. This assessment allows us to either include or exclude any possible dysfunction causing the patient's symptoms. The first step of the algorithm is to analyze this tendinopathy. A tendinous pathology is the most common cause of persistent symptoms in these patients. Pain reproduction through palpation of the epicondylar fossa may indicate the contribution of the extensor tendons to the clinical manifestations³. However, as we will see later on, this tendinous dysfunction may coexist with some other conditions that the clinician must examine during the diagnostic process⁸⁻¹⁷ with the help of the seven-step algorithm. For instance, Myofascial Pain Syndrome or Myofascial Trigger Points of the muscles of the forearm and elbow can increase sensitivity to palpation of the epicondylar region¹⁸, misleading us into believing that the patient has 'epicondylitis' when performing this test.

Moreover, different tests for the diagnosis of epicondylalgia have been typically described: Cozen's test, Mill's test, Polk's test, reduction in grip strength (between 5 and 10%) when using a dynamometer with the elbow extended or the middle finger resistance test (also known as Maudley's test). Despite being recommended by scientific evidence¹⁰⁻²⁰, these tests are insufficient for the clinician since they are positive in many other elbow pathologies apart from a tendinous pathology^{10,13,16,21}.

On the other hand, regarding imaging diagnosis, the presence of anatomo-pathological changes in the extensor tendons of the wrist does not justify that they produce the symptoms⁷ so the clinician must be very careful when interpreting these findings.

Once the clinician has determined the relevance of the dysfunction of the extensor tendons in the patient's symptoms when reproducing pain through palpation of the epicondylar fossa, the second step of the algorithm can be taken ("confirming or excluding radiohumeral synovial plica syndrome").

Second step: confirming or excluding radiohumeral synovial plica syndrome

It is estimated that around 40% of patients with *intractable lateral epicondylitis* present with radiohumeral synovial plica syndrome²². A symptomatic synovial plica is also responsible for the symptoms in other joints but many clinicians are not familiar with its implication in elbow pathologies because there are not many studies about it and the existing reports so far do not seem to reach an agreement. Radiohumeral synovial plica syndrome can exist on its own or together with epicondylitis or other conditions affecting the lateral region of the elbow in a single

patient^{13,22} and therefore its inclusion in the differential diagnosis is key in order to make a correct clinical interpretation. Radiohumeral synovial plicae are remnants of normal embryo development of the articular synovial membrane. We can find one or several of these folds in different sizes, sites, texture and with a different histology. These folds project into the radiohumeral space contiguous with the capsule-ligament complex, found proximal to the edge of the annular ligament within the intra-articular space^{13,23}. It is known that radiohumeral plicae can become symptomatic after a traumatism or due to overuse, although in many cases symptoms appear spontaneously for no clear reason. The symptoms that elbow plicae can develop are multiple and variable and can appear at any age; being focal pain at the posterolateral region of the radiohumeral interline the most characteristic symptom. This pain appears spontaneously in an acute manner accompanying elbow extension movements and with direct palpation during examination. Unfortunately there is currently no gold standard for its diagnosis. However, when assessing the patient, if the clinician sees there is limited range of motion, some blockage in the elbow, snapping, pain during active and/or passive movements with and/or without over pressure, pain when palpating the radiohumeral interline or around the radial head and even some mass in the radial head felt at palpation, radiohumeral synovial plica syndrome must be suspected¹³.

In addition, the use of ultrasounds and MRI, in a compatible clinical context, provide images suggestive of this diagnosis such as: the presence of effusion in the radiohumeral joint, subchondral geodes in the humeral head and/or capitulum and/or luxation or subluxation of the radiohumeral plica. At the same time, these findings allow us to exclude the presence of other intra-articular injuries such as osteochondritis dissecans of the capitulum, allowing the clinician to include radiohumeral synovial plica syndrome as the responsible for the patient's symptoms^{13,23}.

Third step: examining the radial nerve

The third step in the algorithm is the examination of the radial nerve. Radial nerve compression neuropathy shares some clinical characteristics with epicondylitis and can coexist in the same patient so they can be easily confused^{8,10,21,24}. Radial tunnel syndrome¹⁰ and its version with motor deficit, posterior interosseous nerve syndrome^{12,25}, are the most common clinical presentations of damage to the radial nerve in the elbow. Those patients with severe damage of the nerve are easily identified through objective motor dysfunction in the innervated extensor muscles, which commonly affects the common finger extensors in an irregular way²⁶ whereas the majority of cases have a painful syndrome with no sensitivity impairment as clinical presentation since it is a pure motor branch^{8,11,21,24}. If these findings are confirmed, the clinician must refer the patient to a specialist since surgical treatment may be necessary in order to avoid permanent damage¹¹. In most cases the electrophysiological study is normal²¹ but MRI tends to show significant findings¹⁰.

That is why the clinician must suspect this diagnosis when finding the following clinical characteristics: if the patient complains of referred pain in the proximal third of

the dorsal aspect of the forearm^{10,11}, pain at rest, constant pain, radiated pain towards the proximal aspect and/or deep referred pain in the wrist⁷. In addition, some significant findings recorded during the physical examination which are characteristic of the dysfunction of the radial nerve can also arouse the clinician's suspicion, for example: reproduction of symptoms or hypersensitivity to palpation of the radial tunnel, painful palpation at 4–7 cm distal to the lateral epicondyle¹⁰, symptoms during supination counter-resistance²¹ and/or when doing the neurodynamic test of the radial nerve^{7,27}. Ultrasound-guided palpation on the radial nerve projection along the groove of the radial nerve of the humerus can also reproduce the patient's symptoms as well as indicate the presence of some kind of morphological alteration of the radial nerve at the level of the elbow or the presence of a neuroma²⁸ (seen on ultrasounds and/or MRI). These anatomic-pathological findings can add diagnostic value to clinical findings and help the clinician to include the radial nerve as responsible for the patient's symptoms.

Together with these two, the clinician must also consider the posterior antebrachial cutaneous nerve as a possible responsible for these symptoms. This is one of the three nerves in charge of providing cutaneous sensitivity to the forearm with a variable area of cutaneous innervation. The posterior antebrachial cutaneous nerve originates in the radial nerve, between 11 and 18.5 cm proximal to the lateral epicondyle and with a diameter of around 1.9 mm in its origin. This nerve passes through the lateral intermuscular septum before becoming superficial at 6.5–10 cm proximal to the lateral epicondyle²⁹. Its anatomical variation must be taken into account since there are frequently from one to three terminal branches whose size varies in each case and with a variable distal pathway³⁰. In 21% of the cadavers examined we (researchers) found one or two longitudinal branches that are located on average 2.8 cm away from and anterior to the lateral epicondyle and that in most cases (93%) extend along the interval between the brachioradialis and the extensor carpi radialis longus muscles in the proximal aspect of the forearm. In 32% of the specimens examined, the posterior antebrachial cutaneous nerve had a smaller proximal branch and in 86% of the specimens, there was a branch posterior to the lateral epicondyle³¹.

A iatrogenic injury of this nerve is the responsible for the majority of known cases³¹. Although its incidence in *intractable lateral epicondylitis* is still unknown, there is evidence suggesting that the dysfunction of the posterior antebrachial cutaneous nerve may be relevant in cases of persistent lateral elbow pain^{30,32}, which justifies its inclusion in the differential diagnostic process. Any alteration in cutaneous sensitivity in its area of cutaneous innervation will alert the clinician of its implication in the patient's symptoms. If that is the case, it is advisable to perform an anaesthetic block of these branches through ultrasound visualization in order to confirm the diagnosis and assess whether its eventual treatment may be effective^{29,30}.

Fourth step: assessment of articular elbow pathology

In the fourth step of the algorithm the clinician must examine if there is any articular pathology that could account for

the symptoms experienced by the patient with *intractable lateral epicondylitis*. At this stage, the clinician will have already confirmed or ruled out a tendinopathy of the extensor muscles, the radio-humeral plica and the radial nerve as the causes of the patient's problems. The symptoms of snapping, blockage and/or apprehension can coexist with pain typically felt when loading the extensor muscles of the wrist, which commonly appears in epicondylalgia^{3,7}. If the cause of *intractable lateral epicondylitis* is exclusively a tendon and/or neural pathology, these symptoms are not present and therefore if any of them are experienced, the clinician must suspect the presence of an articular dysfunction in the elbow^{13,33}. In these cases, it is advisable to exclude the pathologies in [Table 1](#) through the patient's physical examination and the use of imaging tests.

Injuries to the radial collateral ligament and the lateral ulnar collateral ligament are frequent in patients with chronic epicondylalgia^{8,14}. Significant elbow instability can be found with the use of the conventional physical exams detailed in [Table 1](#). However, there may exist a type of subtle varus instability caused by an injury to the lateral elbow collateral ligament complex, which can be hard to detect in a conventional physical examination due to muscular restrictions. Research on elbow microinstability is scarce but there is evidence to support the inclusion of this clinical syndrome in the differential diagnosis of *intractable lateral epicondylitis*¹⁴. The clinician can rule out this subtle varus instability if no injury to the lateral ligaments is found in the imaging tests (with a negative predictive value of 98.7%). In contrast, if an abnormality is found in these structures, there is the possibility that the patient may have this microinstability. In a study by Kwak et al., 15 out of the 28 abnormal elbow MRIs examined presented subtle instability (with a positive predictive value of 53.6%). In consequence, in order to confirm this articular dysfunction, visualization of the radiocapitellar joint with the use of fluoroscopy is recommended in order to find a widening larger than 1.5 mm at the joint when stressed under anaesthetic¹⁴. Additionally, it is important to underscore that these cases of frank or subclinical instability tend to compensate at muscular level causing Myofascial Pain Syndrome and Myofascial Trigger Points. These symptoms of muscular origin overlap with pain caused by the articular dysfunction, making the diagnosis even harder, and therefore muscular dysfunction must always be assessed when the clinician suspects any alteration in articular stability, as described in step five of the algorithm.

Moreover, in this fourth step the clinician must also exclude the presence of any injury to the annular ligament^{33,35} and of any intra-articular loose bodies in the elbow^{33,34}, as shown in [Table 1](#).

Fifth step: confirming or excluding Myofascial Pain Syndrome and Myofascial Trigger Points

It is known that muscular dysfunction is involved in the chronic symptoms of patients with epicondylalgia¹⁶. In step five of the algorithm, the clinician will either exclude or confirm the presence of myofascial pain. The prevalence of Myofascial Pain Syndrome in lateral elbow pain is variable with a level of implication of the different muscles that

Table 1 Differential diagnosis of *intractable lateral epicondylitis* if the patient has snapping, blockage and/or apprehension.

Condition	Significant characteristics	Test ¹⁹⁻²⁰	Considerations about imaging tests
Posterolateral rotary instability ³⁶⁻³⁸	<p>It is the most common type of elbow instability and its diagnosis is mainly clinical³⁶⁻³⁸. In the physical exam, apprehension tends to be more obvious than subluxation or dislocation due to the patient's pain and protection³⁷.</p> <p>The most common cause is a traumatism but it can also occur iatrogenically due to multiple corticosteroid injections to treat epicondylitis and lateral elbow surgery when inadequately repairing the lateral ulnar collateral ligament or the extensor tendons³⁷.</p>	<p><i>Table-top relocation test.</i></p> <p><i>Stand-up test/chair push-up test.</i></p> <p><i>Push-up test.</i></p> <p><i>Lateral pivot shift test (awake/under anaesthesia)/posterolateral rotatory apprehension test)</i> This is a highly specific test but it has a low degree of sensitivity with the patient awake⁷⁻³⁸.</p> <p><i>Posterolateral drawer test.</i></p>	<p>Many patients have normal or subtly abnormal X-rays³⁷.</p> <p>Although MRIs can be used, it must be borne in mind that injury to the lateral ulnar collateral ligament is not always easily identifiable in cases of chronic posterolateral rotary instability³⁶.</p> <p>There is currently no consensus on the role of this test in the diagnosis of this ligament injury³⁶.</p> <p>Dynamic fluoroscopy and ultrasounds can be useful if the diagnosis is confusing since they show ulnar head subluxation or ulnohumeral widening when either a posterolateral rotary drawer or a supination force on the elbow is applied³⁴.</p>
Varus posteromedial instability ³⁶	<p>It is caused by an injury to the lateral collateral ligament usually after a traumatism which can also produce a fracture of the anteromedial aspect or coronoid apophysis³⁶.</p>	<p><i>Gravity-assisted varus stress test</i>³⁶.</p>	<p>The elbow varus stress test under anaesthetics is the diagnostic gold standard, an opening occurs in the ulnohumeral region³⁶.</p> <p>There is currently no consensus on the role of ultrasounds in the assessment of the lateral ligament complex in relation to this instability³⁶.</p>
Valgus instability ^{36,39}	<p>Also known as "medial collateral ligament insufficiency".</p> <p>This ligament can be damaged due to traumatism or repetitive elbow overuse and it is commonly found in athletes participating in overhead throwing sports³⁹.</p>	<p><i>Moving valgus stress test.</i></p> <p><i>Valgus stress test/ligamentous instability test.</i></p> <p><i>Milking manoeuvre.</i></p>	<p>Arthrography with a saline or gadolinium-enhanced MRI has a sensitivity of 97% and a specificity of 100%³⁶.</p> <p>Valgus stress X-rays can be useful³⁹, as well as dynamic ultrasounds. They both can show medial joint instability when valgus stress is applied, assessing the medial ulnohumeral articular space. Additionally, the continuity of the ulnar collateral ligament can be observed with ultrasounds⁴⁰. However, it must be borne in mind that in both tests an articular widening is observed which is expected in the dominant limb of overhead athletes, making outcome interpretation and diagnosis more difficult³⁶.</p>

Table 1 (Continued)

Condition	Significant characteristics	Test ¹⁹⁻²⁰	Considerations about imaging tests
Annular ligament injury ³³⁻³⁵	<p>Some of these patients do not experience snapping or blockage.</p> <p>Patients with this injury can have a history of ulnar head fractures/luxation or malformations, fractures in the proximal radioulnar area, distal humeral fractures or prior elbow arthroscopy.</p> <p>It is possible to find instability in the annular ligament together with instability in the radioulnar joint³⁵.</p>	Symptoms may be caused by the gliding of the annular ligament on the ulnohumeral joint when bending and stretching the elbow ³⁵ .	MRI or dynamic ultrasounds can help confirm the diagnosis ³⁵ .

Table 2 Muscles whose Myofascial Trigger Points can be involved in *intractable lateral epicondylitis* and that the clinician must assess (from highest to lowest prevalence).

- Extensor carpi radialis brevis muscle
 - Finger extensor muscle
 - Extensor carpi radialis longus muscle
- Brachioradial muscle
- Brachial triceps (mainly lateral fibres of the medial head)
- Supinator muscle
- Anconeus muscle
- Extensor carpi ulnaris muscle

varies according to different studies⁴¹⁻⁴⁴. Although subjective and manual physical examination can be confusing, it is recommended that the clinician includes the following muscles in the differential diagnosis (Table 2).

Pain reproduction through the precise and analytical palpation of each muscle and/or the reproduction of symptoms with the use of dry needling can help the clinician to confirm this dysfunction as an element responsible for the symptoms¹⁶. The examination of each muscle under ultrasound control greatly improves the reliability of the assessment.

Sixth step: excluding the cervical spine

In the sixth step of the algorithm referred pain from the cervical spine must be excluded as the cause of chronic elbow pain⁷. In order to do this the clinician must answer the following questions: is it the neck or the elbow the region that reproduces the symptoms? Is there radicular pain? During the assessment, the clinician will ask the patient about the presence of concomitant neck pain and will assess the presence of any restriction in active range of motion of the neck and any neurological symptoms in the upper limb. During the

patient's physical exam, the reproduction of lateral elbow pain through manual palpation and/or active and/or passive movements of the cervical or thoracic spine⁴⁴⁻⁴⁶ must alert the clinician as they can indicate the involvement of the spine in the patient's *intractable lateral epicondylitis*^{47,48}.

Seventh step: examining central sensitization and psychosocial factors

Central sensitization is involved in the pathophysiology of extensor tendon epicondylitis just as in other tendinopathies of the upper limb and in many musculoskeletal pain syndromes^{7,49} so its assessment will be the focus of the seventh and last step in this differential diagnosis algorithm. Anxiety and depression can also be frequently present in these patients^{17,50}. Although further research is needed, the clinician is advised to consider psychosocial factors and include psychological assessment tools when dealing with patients with *intractable lateral epicondylitis* if necessary, once the rest of diagnoses have been confirmed or excluded.

Discussion

Epicondylalgia is the most common cause of lateral elbow pain in adults and when treatments do not seem to work, it is labelled "*intractable lateral epicondylitis*". Does this clinical entity really exist or is it a misdiagnosis that consequently results in wrong treatment? The amount of structures that can cause the symptoms is really big and several clinical syndromes can coexist in lateral elbow pain. Moreover, the traditional tests used to diagnose epicondylalgia do not help in the screening of the different pathologies because many of them are positive and they are probably more useful as functional assessment parameters than as differential diagnostic tools. On the other hand, imaging tests alone are not enough for the diagnosis since most of the structures responsible for the symptoms in *intractable lateral epicondylitis* can be asymptomatic in spite of the fact

that test images can show anatomic-pathological changes or the other way round: they can have a normal appearance and be symptomatic. That is why the clinician must be very meticulous when checking if there is any correlation between tissue changes and the patient's subjective and physical findings, which explains why a comprehensive and precise physical exam of the patient is essential and decisive for a correct clinical interpretation.

On the other hand, it must be taken into account that emotional and psychological factors can play a role in the perpetuation of symptoms. Nevertheless, there is the possibility that these are the consequence and not "the cause" of the pain and disability as it is normally claimed. For this reason the clinician must avoid attributing responsibility for pain chronification in patients with epicondylalgia to central sensitization and to psychosocial factors before making a comprehensive differential diagnosis of the different clinical syndromes that can coexist and get confused thus avoiding erroneous interpretations in the management of these patients⁵¹.

Conclusion

Intractable lateral epicondylitis is a complex entity comprising different clinical syndromes that can coexist and get mixed up. That is why it is necessary to perform a systematic and comprehensive assessment of the patient like the one proposed in this article, where the musculotendinous articular and neural dysfunction responsible for the patient's symptoms is rigorously analyzed from highest to lowest prevalence. This seven-step differential diagnostic algorithm can be very useful for those clinicians dealing with these patients and can be base for future research, focusing on the identification of different subgroups of patients with *intractable lateral epicondylitis* for a correct clinical interpretation.

Conflict of interest

The authors declare that there is no conflict of interest.

References

- Luk JK, Tsang RC, Leung HB. Lateral epicondylalgia: midlife crisis of a tendon. *Hong Kong Med J*. 2014;20:145–51, <http://dx.doi.org/10.12809/hkmj134110> [Epub 28.02.14; PMID: 24584568].
- Oh DS, Kang TH, Kim HJ. Pulsed radiofrequency on radial nerve under ultrasound guidance for treatment of intractable lateral epicondylitis. *J Anesth*. 2016;30:498–502, <http://dx.doi.org/10.1007/s00540-016-2146-9> [Epub 20.02.16; PMID: 26896944].
- Vaquero-Picado A, Barco R, Antuña SA. Lateral epicondylitis of the elbow. *EFORT Open Rev*. 2017;1:391–7, <http://dx.doi.org/10.1302/2058-5241.1.000049> [PMID: 28461918; PMID: PMC5367546].
- Keijsers R, de Vos RJ, Kuijer PPF, van den Bekerom MP, van der Woude HJ, Eygendaal D. Tennis elbow. *Shoulder Elbow*. 2019;11:384–92, <http://dx.doi.org/10.1177/1758573218797973> [Epub 18.08.18; PMID: 31534489; PMID: PMC6739751].
- Cutts S, Gangoo S, Modi N, Pasapula C. Tennis elbow: a clinical review article. *J Orthop*. 2019;17:203–7, <http://dx.doi.org/10.1016/j.jor.2019.08.005> [PMID: 31889742; PMID: PMC6926298].
- Lai WC, Erickson BJ, Mlynarek RA, Wang D. Chronic lateral epicondylitis: challenges and solutions. *Open Access J Sports Med*. 2018;9:243–51, <http://dx.doi.org/10.2147/OAJSM.S160974> [PMID: 30464656; PMID: PMC6214594].
- Coombes BK, Bisset L, Vicenzino B. Management of lateral elbow tendinopathy: one size does not fit all. *J Orthop Sports Phys Ther*. 2015;45:938–49, <http://dx.doi.org/10.2519/jospt.2015.5841> [Epub 17.09.15].
- Kotnis NA, Chiavaras MM, Harish S. Lateral epicondylitis and beyond: imaging of lateral elbow pain with clinical–radiologic correlation. *Skeletal Radiol*. 2012;41:369–86.
- Qi L, Zhu ZF, Li F, Wang RF. MR imaging of patients with lateral epicondylitis of the elbow: is the common extensor tendon an isolated lesion? *PLOS ONE*. 2013;8:e79498, <http://dx.doi.org/10.1371/journal.pone.0079498> [eCollection 2013].
- Moradi A, Ebrahimzadeh MH, Jupiter BJB. Radial tunnel syndrome diagnostic and treatment dilemma. *Arch Bone Jt Surg*. 2015;3:156–62.
- Carter GT, Weiss MD. Diagnosis and treatment of work-related proximal median and radial nerve entrapment. *Phys Med Rehabil Clin N Am*. 2015;26:539–49.
- Cha YK, Kim SJ, Park NH, Kim JY, Kim JH, Park JY. Magnetic resonance imaging of patients with lateral epicondylitis: relationship between pain and severity of imaging features in elbow joints. *Acta Orthop Traumatol Turc*. 2019;53:366–71, <http://dx.doi.org/10.1016/j.aott.2019.04.006> [Epub 28.04.19].
- Aguililla Liñan JM, Miguel Pérez MI, Palau González J, Möller Parera I, Martinoli C. A comprehensive review of radiohumeral synovial plicae for a correct clinical interpretation in intractable lateral epicondylitis. *Curr Rev Musculoskelet Med*. 2020;13:385–90, <http://dx.doi.org/10.1007/s12178-020-09636-w> [PMID: 32458355; PMID: PMC7340713].
- Kwak SH, Lee SJ, Jeong HS, Do MU, Suh KT. Subtle elbow instability associated with lateral epicondylitis. *BMC Musculoskelet Disord*. 2018;19:136, <http://dx.doi.org/10.1186/s12891-018-2069-8> [PMID: 29734945; PMID: PMC5938800].
- Coel M, Yamada CY, Ko J. MR imaging of patients with lateral epicondylitis of the elbow (tennis elbow): importance of increased signal of the anconeus muscle. *AJR Am J Roentgenol*. 1993;161:1019–21.
- Shmushkevich Y, Kalichman L. Myofascial pain in lateral epicondylalgia: a review. *J Bodyw Mov Ther*. 2013;17:434–9, <http://dx.doi.org/10.1016/j.jbmt.2013.02.003> [Epub 21.04.13].
- Aben A, De Wilde L, Hollevoet N, Henriquez C, Vandeweerdt M, Ponnet K, et al. Tennis elbow: associated psychological factors. *J Shoulder Elbow Surg*. 2018;27:387–92, <http://dx.doi.org/10.1016/j.jse.2017.11.033>.
- Hong CZ. New trends in myofascial pain syndrome. *Zhonghua Yi Xue Za Zhi (Taipei)*. 2002;65:501–12 [PMID: 12583512].
- Smith MV, Lamplot JD, Wright RW, Brophy RH. Comprehensive review of the elbow physical examination. *J Am Acad Orthop Surg*. 2018;26:678–87, <http://dx.doi.org/10.5435/JAAOS-D-16-00622> [PMID: 30095513].
- Zwerus EL, Somford MP, Maissan F, Heisen J, Eygendaal D, van den Bekerom MP. Physical examination of the elbow, what is the evidence? A systematic literature review. *Br J Sports Med*. 2018;52:1253–60, <http://dx.doi.org/10.1136/bjsports-2016-096712> [Epub 01.03.17; PMID: 28249855].

21. Bouche P. Compression and entrapment neuropathies. *Handb Clin Neurol.* 2013;115:311–66.
22. Kongmalai P, Chanlalit C. Demographic causes of chronic lateral elbow pain along arthroscopic criteria. *J Med Assoc Thai.* 2016;99 Suppl. 8:S79–83 [PMID: 29901918].
23. Wong JS, Lalam R. Plicae: where do they come from and when are they relevant? *Semin Musculoskelet Radiol.* 2019;23:547–68, <http://dx.doi.org/10.1055/s-0039-1693979> [Epub 25.09.19; PMID: 31556089].
24. Gürçay E, Karahmet ÖZ, Kara M, Onat SS, Ata AM, Ünlü E, et al. Ultrasonographic evaluation of the radial nerves in patients with unilateral refractory lateral epicondylitis. *Pain Med.* 2017;18:396–402, <http://dx.doi.org/10.1093/pm/pnw181> [PMID: 27477582].
25. Anania P, Fiaschi P, Ceraudo M, Balestrino A, Zaotini F, Martinoli C, et al. Posterior interosseous nerve entrapments: review of the literatura. Is the entrapment distal to the arcade of Frohse a really rare condition? *Acta Neurochir (Wien).* 2018;160:1857–64, <http://dx.doi.org/10.1007/s00701-018-3615-8> [Epub 04.08.18; PMID: 29974240].
26. Latef TJ, Bilal M, Vetter M, Iwanaga J, Oskouian RJ, Tubbs RS. Injury of the radial nerve in the arm: a review. *Cureus.* 2018;10:e2199, <http://dx.doi.org/10.7759/cureus.2199> [PMID: 29666777; PMCID: PMC5902095].
27. Koulidis K, Veremis Y, Anderson C, Heneghan NR. Diagnostic accuracy of upper limb neurodynamic tests for the assessment of peripheral neuropathic pain: a systematic review. *Musculoskelet Sci Pract.* 2019;40:21–33, <http://dx.doi.org/10.1016/j.msksp.2019.01.001> [Epub 12.01.19; PMID: 30665045].
28. Shi M, Qi H, Ding H, Chen F, Xin Z, Zhao Q, et al. Electrophysiological examination and high frequency ultrasonography for diagnosis of radial nerve torsion and compression. *Medicine (Baltimore).* 2018;97:e9587, <http://dx.doi.org/10.1097/MD.00000000000009587> [PMID: 29480857; PMCID: PMC5943862].
29. Finneran JJ, Sandhu N. Ultrasound-guided posterior antebrachial cutaneous nerve block: technical description and block distribution in healthy volunteers. *J Ultrasound Med.* 2019;38:239–42, <http://dx.doi.org/10.1002/jum.14678> [Epub 06.05.18; PMID: 29732596].
30. Rose NE, Forman SK, Dellon AL. Denervation of the lateral humeral epicondyle for treatment of chronic lateral epicondylitis. *J Hand Surg Am.* 2013;38:344–9, <http://dx.doi.org/10.1016/j.jhsa.2012.10.033> [PMID: 23351911].
31. Starr BW, Lee DS, Stern PJ. Anatomy of the posterior antebrachial cutaneous nerve, revisited. *J Hand Surg Am.* 2020;45:360.e1–4, <http://dx.doi.org/10.1016/j.jhsa.2019.08.011> [Epub 22.10.19; PMID: 31653469].
32. Berry N, Neumeister MW, Russell RC, Dellon AL. Epicondylectomy versus denervation for lateral humeral epicondylitis. *Hand (N Y).* 2011;6:174–8, <http://dx.doi.org/10.1007/s11552-011-9318-8> [Epub 25.02.11; PMID: 22654700; PMCID: PMC3092894].
33. Feller RJ, Gil JA, DaSilva M. Snapping at the lateral aspect of the elbow: a case report and review of the literature. *JBJS Case Connect.* 2018;8:e48, <http://dx.doi.org/10.2106/JBJS.CC.17.00198> [PMID: 29995662].
34. Camp CL, Smith J, O'Driscoll SW. Posterolateral rotatory instability of the elbow: Part II. Supplementary examination and dynamic imaging techniques. *Arthrosc Tech.* 2017;6:e407–11, <http://dx.doi.org/10.1016/j.eats.2016.10.012> [PMID: 28580260; PMCID: PMC5442982].
35. Kerver N, Boedda AV, Gerritsma-Bleeker CLE, Eygen-daal D. Snapping of the annular ligament: a uncommon injury characterised by snapping or locking of the elbow with good surgical outcomes. *Knee Surg Sports Traumatol Arthrosc.* 2019;27:326–33, <http://dx.doi.org/10.1007/s00167-018-5076-2> [Epub 02.08.18; PMID: 30073382].
36. Karbach LE, Elfar J. Elbow instability: anatomy, biomechanics diagnostic maneuvers, and testing. *J Hand Surg Am.* 2017;42:118–26, <http://dx.doi.org/10.1016/j.jhsa.2016.11.025> [PMID: 28160902; PMCID: PMC5821063].
37. Fedorka CJ, Oh LS. Posterolateral rotatory instability of the elbow. *Curr Rev Musculoskelet Med.* 2016;9:240–6, <http://dx.doi.org/10.1007/s12178-016-9345-8> [PMID: 27194295; PMCID: PMC4896886].
38. Conti Mica M, Caekebeke P, van Riet R. Lateral collateral ligament injuries of the elbow – chronic posterolateral rotatory instability (PLRI). *EFORT Open Rev.* 2017;1:461–8, <http://dx.doi.org/10.1302/2058-5241.160033> [PMID: 28461924; PMCID: PMC5367576].
39. Willemot L, Hendriks FR, Byrne AM, van Riet RP. Valgus instability of the elbow: acute and chronic form. *Obere Extremit.* 2018;13:173–9, <http://dx.doi.org/10.1007/s11678-018-0465-1> [Epub 30.05.18; PMID: 30220920; PMCID: PMC6132395].
40. Hendawi TK, Rendos NK, Warrell CS, Hackel JG, Jordan SE, Andrews JR, et al. Medial elbow stability assessment after ultrasound-guided ulnar collateral ligament transection in a cadaveric model: ultrasound versus stress radiography. *J Shoulder Elbow Surg.* 2019;28:1154–8, <http://dx.doi.org/10.1016/j.jse.2018.11.060> [Epub 13.02.19; PMID: 30770313].
41. Fernández-Carnero J, Fernández-de-Las-Peñas C, de la Llave-Rincón AI, Ge HY, Arendt-Nielsen L. Prevalence of and referred pain from myofascial trigger points in the forearm muscles in patients with lateral epicondylalgia. *Clin J Pain.* 2007;23:353–60, <http://dx.doi.org/10.1097/AJP.0b013e318033785> [PMID: 17449997].
42. Fernández-Carnero J, Fernández-de-las-Peñas C, de la Llave-Rincón AI, Ge HY, Arendt-Nielsen L. Bilateral myofascial trigger points in the forearm muscles in patients with chronic unilateral lateral epicondylalgia: a blinded, controlled study. *Clin J Pain.* 2008;24:802–7, <http://dx.doi.org/10.1097/AJP.0b013e31817cb79> [PMID: 18936598].
43. Fernández-Carnero J, Ge HY, Kimura Y, Fernández-de-Las-Peñas C, Arendt-Nielsen L. Increased spontaneous electrical activity at a latent myofascial trigger point after nociceptive stimulation of another latent trigger point. *Clin J Pain.* 2010;26:138–43, <http://dx.doi.org/10.1097/AJP.0b013e3181bad736> [PMID: 20090441].
44. Berglund KM, Persson BH, Denison E. Prevalence of pain and dysfunction in the cervical and thoracic spine in persons with and without lateral elbow pain. *Man Ther.* 2008;13:295–9, <http://dx.doi.org/10.1016/j.math.2007.01.015> [Epub 17.10.07; PMID: 17942362].
45. Coombes BK, Bisset L, Vicenzino B. Bilateral cervical dysfunction in patients with unilateral lateral epicondylalgia without concomitant cervical or upper limb symptoms: a cross-sectional case–control study. *J Manipulative Physiol Ther.* 2014;37:79–86, <http://dx.doi.org/10.1016/j.jmpt.2013.12.005> [Epub 28.12.13; PMID: 24378321].
46. Thoomes EJ, van Geest S, van der Windt DA, Falla D, Verhagen AP, Koes BW, et al. Value of physical tests in diagnosing cervical radiculopathy: a systematic review. *Spine J.* 2018;18:179–89, <http://dx.doi.org/10.1016/j.spinee.2017.08.241> [Epub 31.08.17; PMID: 28838857].
47. Cook C, Learman K, Showalter C, O'Halloran B. The relationship between chief complaint and comparable sign in patients with spinal pain: an exploratory study. *Man Ther.* 2015;20:451–5,

- <http://dx.doi.org/10.1016/j.math.2014.11.007> [Epub 22.11.14; PMID: 25498410].
48. Cook C, Brown C, Isaacs R, Roman M, Davis S, Richardson W. Clustered clinical findings for diagnosis of cervical spine myelopathy. *J Man Manip Ther.* 2010;18:175–80, <http://dx.doi.org/10.1179/106698110X12804993427045> [PMID: 22131790; PMCID: PMC3113267].
49. Plinsinga ML, Brink MS, Vicenzino B, van Wilgen CP. Evidence of nervous system sensitization in commonly presenting and persistent painful tendinopathies: a systematic review. *J Orthop Sports Phys Ther.* 2015;45:864–75, <http://dx.doi.org/10.2519/jospt.2015.5895> [Epub 21.09.15; PMID: 26390275].
50. Gürçay E, Tamkan AU, Karaahmet ÖZ, Tombak Y, Güzel Ş, Çakci A. Depression and somatization in refractory lateral epicondylitis. *Arch Rheumatol.* 2019;34:367–70, <http://dx.doi.org/10.5606/ArchRheumatol.2019.7139> [PMID: 32010884; PMCID: PMC6974392].
51. Rhyou IH, Kim KW. Is posterior synovial plica excision necessary for refractory lateral epicondylitis of the elbow? *Clin Orthop Relat Res.* 2013;471:284–90, <http://dx.doi.org/10.1007/s11999-012-2585-z> [Epub 11.08.12; PMID: 22965262; PMCID: PMC3528936].